## REMOVAL SITE EVALUATION OF MERCURY IN SOIL AND GROUNDWATER AT FORMER MERCURY FULMINATE FACILITY

#### RICHMOND FIELD STATION

Richmond, California

May 21,1990

#### Submitted to:

University of California Richmond Field Station Richmond, California

#### Prepared by:

Jonas & Associates Inc.
Environmental Consultants

1056 Dale Place
Concord, California
(415) 676-8554

## TABLE OF CONTENTS

				PAGE
LIST	OF TABLES			iv
LIST	OF FIGURES		•	v
LIST	OF APPENDIC	ES		vi
EXE	CUTIVE SUMM	ARY		vii
1.0	INTRODUCT	ION		1
	1.1 Site Loca	tion		4
	1.2Site Statu	ıs		4
	1.3 Chronolog	gy of Ever	nts	8
2.0	INITIAL SITE	EVALUA	ATION	11
	2.1 Site Desc	ription		11
	2.1.1	Property	History and Adjacent Land Use	11
			Property History Adjacent Land Use	11 11
	2.1.2	Environ	nental Setting	13
		2.1.2.2 2.1.2.3	Regional Geology Local Geology Surface Water Regional Hydrogeology and Groundwater Quality	13 13 14
			2.1.2.4.1 Regional Hydrogeology 2.1.2.4.2 Local Hydrogeology 2.1.2.4.3 Groundwater Quality 2.1.2.4.4 Surface Water Quality	14 14 16 18

## TABLE OF CONTENTS (Continued)

			<u>PAGE</u>
	2.2 Summary	of the Previous Studies (1981 - 1988)	19
	<b>2.2.</b> 1	Previous Investigations	19
3.0	CURRENT S	OIL SAMPLING STUDY CONDUCTED BY	22
	3.1 Sampling	Methods and Procedures	22
		Sample Collection and Analysis 3.1.1.1 Task 1 - Soil Sample Collection from 0 to 3	22
	/	Feet Depth Range	22
		3.1.1.2 Task 2 - Surface Soil Sample Collection 3.1.1.3 Task 3 - Collection of Soil Sample from the	24
		Groundwater Monitoring Well (MW-1) Borehole	24
	3.1.2	Decontamination and Post-Sampling Procedures	25
		Sample Documentation	26
		Sample Identification Numbers	26
		Field Logbooks	26
		Sample Labels	27
		Chain-of-Custody Records	27 28
		Quality Control Samples	28 28
	3.1.9	Sample Shipment	28
	3.2 Discussion	on of Results of Soil Analyses	28
		Results of Analyses for Mercury in Soil Samples	<del>28</del>
	3.2.2	Results of Analyses for Sulfate and Nitrate in Soil Samples	38
	3.2.3	Results of Analyses for Copper and Zinc in Soil Samples	39
	3.2.4	Results of EP Toxicity and Waste Extraction (WET) Tests fo Mercury in Soil Samples	r 39
4.0		ATER MONITORING WELL DRILLING, CONSTRUCTION, ENT, AND SAMPLING	40
	4.1 Drilling 1	Methods	40
	4.1 Drilling 1		44
	4.3 Well Dev		45
		Collection Procedures and Analysis	45

#### TABLE OF CONTENTS (Continued)

		PAGE
	4.5 Results of Analyses for Mercury and Total Dissolved Solids (TDS) in Sample	Water 46
5.0	THE CHEMISTRY, TOXICOLOGY, AND MIGRATION PATHWAYS OF MERCURY AND MERCURY FULMINATE	46
	5.1 Chemistry of Mercury and Mercury Fulminate	46
	5.1.1 Chemistry of Mercury 5.1.2 Mercury Fulminate	46 46
	5.3 Fate and Transport of Mercury	49
	<ul><li>5.3.1 Volatilization</li><li>5.3.2 Sorption</li><li>5.3.3 Biotransformation</li></ul>	49 49 49
	5.3 Toxicology of Mercury	50
6.0	SUMMARY AND RECOMMENDATIONS	51
	6.1 Summary of the Investigation 6.2 Summary of Results 6.3 Recommendations	51 52 52
REFI	ERENCES	54

#### LIST OF TABLES

	<u>PAGE</u>
Borehole Stratigraphy	13
Results of Existing Richmond Field Station (RFS) Well Sample Analyses (EAL Corporation, 1984)	16
Results of ICI Americas Well Sample Analyses	17
Results of Slough and Storm Drain Sample Analyses (EAL Corporation, 1984)	18
Results of 1988 Soil Samples Collected From the Former Mercury Fulminate Facility (CH2M Hill, 1988)	22
Surface Samples and Corresponding Borehole Locations (J & A, 1990)	25
Mercury in 0 - 3 Feet of Soil, Richmond Field Station (J & A, 1989 - 1990)	29
Mercury in Surface Soil, Richmond Field Station (J & A, 1990)	37
Sulfate & Nitrate in 0 - 3 Feet of Soil, Richmond Field Station (J & A, 1990)	38
Copper & Zinc in Surface Soil, Richmond Field Station (J & A, 1990)	39
	Results of Existing Richmond Field Station (RFS) Well Sample Analyses (EAL Corporation, 1984)  Results of ICI Americas Well Sample Analyses  Results of Slough and Storm Drain Sample Analyses (EAL Corporation, 1984)  Results of 1988 Soil Samples Collected From the Former Mercury Fulminate Facility (CH2M Hill, 1988)  Surface Samples and Corresponding Borehole Locations (J & A, 1990)  Mercury in 0 - 3 Feet of Soil, Richmond Field Station (J & A, 1989 - 1990)  Mercury in Surface Soil, Richmond Field Station (J & A, 1990)  Sulfate & Nitrate in 0 - 3 Feet of Soil, Richmond Field Station (J & A, 1990)  Copper & Zinc in Surface Soil,

#### LIST OF FIGURES

		PAGE
Figure 1-1	Regional Map	5
Figure 1-2	Site Map - General Area of Investigation	6
Figure 1-3	Current and Historic Land Use in Study Area	7
Figure 2-1	Proposed Facility	12
Figure 2-2	Previous Sampling Locations	20
Figure 2-3	Previous Soil Mercury Results	21
Figure 3-1	Location of Soil Samples and Monitoring Well (J & A, 1989 - 1990)	23
Figure 3-2	Sample Label	27.
Figure 3-3	Mercury in Soil at the Study Area, 0 - 3 Feet (J & A, 1989 - 90)	30
Figure 3-4	Past and Present Structures and Soil Mercury Concentration	31
Figure 3-5	Elevated Mercury Levels in Soil, 0 - 3 Feet (J & A, 1989 - 90)	32
Figure 3-6	Elevated Mercury Levels in Surface Soil (J & A, 1989 - 90)	33
Figure 3-7	Sulfate & Nitrate in Soil, 0 - 3 Feet (J & A, 1989 - 90)	34
Figure 3-8	Copper in Surface Soil and at 0 - 3 Feet (J & A, 1989 - 90)	35
Figure 3-9	Zinc in Surface Soil and at 0 - 3 Feet (J & A, 1989 - 90)	36
Figure 4-1	MW-1 Construction Details	41
Figure 4-2	MW-1 Stratigraphy and Soil Analytical Results	42
Figure 4-3	MW-A Construction Details	43
Figure 4-4	MW-1, Probable Groundwater Flow, and Elevated Soil Mercury	47

In 1982, DHS collected 17 additional soil samples from the RFS facility and analyzed the samples for a large suite of inorganic constituents. Mercury was not detected in any of the samples collected by DHS in 1982. Two of these samples were from the same locations as the earlier DHS samples collected in 1981. In the 1982 sampling effort, none of the inorganic constituents in soil samples exceeded the Total Threshold Limit Concentrations (TTLC) set by DHS to identify a material as hazardous. DHS concluded that no mercury remained on the RFS property and recommended that RFS should be removed from the list of abandoned hazardous waste sites.

In 1988, CH2M Hill collected additional soil samples at RFS. The purpose of this investigation was to follow-up on previous sampling activities and to supplement and/or clarify analytical results.

During this investigation three soil samples were collected in the general area of the former mercury fulminate facility. All soil samples were analyzed for metals, pesticides and PCBs. Copper and zinc in soil samples were detected below TTLC levels. Mercury in soils exceeded TTLC levels. Pesticides and PCBs were not detected in soil samples collected.

The Phase I investigation, which is the subject of this report, included collection of 147. discrete soil samples (0 - 3 feet) collected from 47 boreholes; analysis of 49 composite soil samples for mercury; analysis of 4 composite soil samples for sulfate, nitrate, copper and zinc; and collection and analysis of 10 surface soil samples (0 - 3 inches) for mercury. Two soil samples were also analyzed for mercury using EP Toxicity Extraction and Waste Extraction (WET) tests. In addition, one groundwater monitoring well was installed and groundwater was sampled and analyzed for mercury. A total of 4 additional composite soil samples were collected from the monitoring well borehole and analyzed for mercury.

A summary of the findings for Phase I indicates that mercury is present in the soil above the TTLC level. No mercury was detected in the groundwater sample analyzed. Copper and zinc were present in the soil below TTLC levels, and sulfate and nitrate were detected in soil in low concentrations. Mercury was detected above the Soluble—Threshold Concentration Level (STLC: 0.2 mg/l) using the Waste Extraction (WET) test. Mercury was not detected above the method detection limit of 0.02 mg/l in the soil sample that was analyzed using the EP Toxicity extraction method.

It is recommended that the following tasks be incorporated into Phase II of this investigation which is intended to serve as a preliminary Public Health and Environmental Evaluation (PHEE).

- Perform a preliminary Public Health and Environmental Evaluation (PHEE);
- Conduct field investigations as necessary to collect any additional data needed for the PHEE;
- · Continue groundwater sampling and analysis; and

• Using information developed by the PHEE provide a quantitative basis for selection of an appropriate removal action.

However, the exact scope of work for the Phase II of the investigation will be finalized after UC's meeting(s) with DHS and other regulatory agencies review of this report

## LIST OF APPENDICES

Appendix A J&A Soil Sampling Laboratory Results and Chain-of-Custody Forms

#### **EXECUTIVE SUMMARY**

The University of California (UC), Richmond Field Station (RFS) is located at 1301 South 46th Street in Richmond, California 94804. Beginning in the 1840s, several explosives manufacturing companies were operating in an area that is now the south-eastern portion of RFS. At the beginning of the 1900s, the California Cap Company (CCC) bought the land and one of the explosive manufacturing operations from the Hercules Powder Company.

CCC ceased operations at the end of World War II in 1945. UC purchased the property in 1950. The property is currently owned and operated by UC. Several different facilities, including research laboratories and administrative offices are located on this property. Currently, RFS is considering an addition of a new building at their facility. Due to the proximity of this proposed building to a former explosives manufacturing facility where mercury fulminate was produced, RFS decided to initiate the current investigation of mercury. Earlier investigations had indicated the presence of mercury in soils near this facility (DHS, 1982; CH2M Hill, 1988). Jonas & Associates Inc. (J & A) was retained in December 1989, under order 0-200248-TR to conduct the current investigation. The mercury contamination investigation is being performed according to the "National Oil and Hazardous Substances Pollution Contingency Plan, February 2, 1990 (NCP)."

Based on NCP requirements, UC has initiated a phased approach for the investigation of mercury contamination. Phase I is the performance of a Removal Site Evaluation, which has now been completed and the results are presented in this report. Following the Removal Site Evaluation, consideration of the need for Removal Action will commence as Phase II of the investigation. If Phase II evaluation indicates that removal action is appropriate, an Engineering Evaluation/Cost Analysis (EE/CA) or its equivalent will be performed. If collection of additional environmental samples becomes necessary, then a sampling and analysis plan will be prepared and submitted for regulatory review.

The scope of work for Phase I of the investigation was focused on the general area of the former mercury fulminate facility. The emphasis was placed on determining mercury concentration in soil. However, during the investigation a preliminary evaluation of nitrate, sulfate, copper, and zinc in soil samples was also conducted. In addition, an evaluation of mercury in groundwater was initiated by installation of one groundwater monitoring well in the study area. Groundwater from this well was analyzed for mercury. Mercury was not detected above the instrument detection limit of 0.001 mg/l in the groundwater sample.

RFS was included on the Department of Health Services (DHS) list of abandoned hazardous waste sites in 1981 because of CCC's explosive manufacturing activities. DHS began an investigation of the RFS property in this same year, at which time samples from five locations were collected and analyzed for mercury. Mercury was detected in two of these samples.

#### 1.0 INTRODUCTION

The University of California at Berkeley (UC) has initiated a focused investigation of potential mercury contamination at the former mercury fulminate facility located within the Richmond Field Station (RFS), Richmond, California. Earlier investigations (summarized in Section 1.3) indicated the presence of mercury in soils within the study area. Jonas & Associates Inc. (J & A) was retained in December 1989, under order 0-200248-TR, to initiate an Investigation of mercury contamination at the former mercury fulminate facility. The mercury contamination investigation is being performed according to the "National Oil and Hazardous Substances Pollution Contingency Plan, February 2, 1990 (NCP)." Based on NCP requirements, UC has initiated a phased approach for the investigation of mercury contamination. These phases are as follow:

Phase I - Performance of a Removal Site Evaluation which includes:

· Performance of a removal preliminary assessment

A removal preliminary assessment may include collection and review of data such as site management practices, information from generators, photographs, analysis of historical photographs, literature searches, and personal interviews conducted as appropriate.

Performance of a removal site inspection

A removal site inspection may be performed if more information is need. Such inspection may include a perimeter (off-site) or on-site inspection, taking into consideration whether such inspection can be performed safely.

Phase I has now been completed and the results are presented in this report.

Phase II - Following the completion of the removal site evaluation, consideration of the need for removal action will commence. Under the NCP, the following factors are to be considered by the lead agency in determining the appropriateness of a removal action:

As described in greater detail in this report, the analysis is approached as a "removal" action for NCP purposes due to the following factors: (1) the mercury contaminated soil appears to be localized and the mercury does not appear to be migrating, (2) no groundwater contamination has been detected, (3) all of the likely response actions are included in the listing of removal actions set forth in the NCP, 40 C.F.R. S 300.415 (d), (4) any response actions in the areas investigated would likely be independent from other potential response actions at the site, and (5) given the anticipated volume of the contaminated soil, no presently anticipated cleanup option would approach or exceed the \$2 million threshold for removal actions.

Containment, treatment, disposal, or incineration of hazardous materials -where needed to reduce the likelihood of human, animal, or food chain
exposure.

This report presents the findings of Phase I of the investigation and describes the site and the site's current status, defines the objectives of the investigation, provides a chronology of events, includes a general information on the geology and hydrology of the site and the region, presents the results of the recent soil and groundwater sampling investigation and provides a general discussion of the chemistry, toxicology, and migration pathways of mercury. A preliminary public health and environmental evaluation, proposed as part of the Phase II of the investigation, would present an indepth analysis of the potential risks associated with exposure to mercury. If the Phase II evaluation indicates that removal action is appropriate, a Phase III EE/CA will be undertaken.

This report, Removal Site Evaluation of Mercury in Soil and Groundwater at Former Mercury Fulminate Facility, Richmond Field Station, is organized into six major sections:

Section 1: Introduction

Section 2: Initial Site Evaluation

Section 3: Current Soil Sampling Study Conducted by Jonas & Associates Inc.

Section 4: Groundwater Monitoring Well Drilling, Construction, Development,

Sampling and Analysis

Section 5: The Chemistry, Toxicology, and Migration Pathways of Mercury and

Mercury Fulminate

Section 6: Summary and Recommendations

Section 1 describes the site and the site's current status, defines the objectives of the investigation, and provides a chronology of the events occurring in the general area of the former mercury fulminate facility.

Section 2 provides a site description, describes the geology and hydrology of the site and region, and presents a summary of previous investigations performed by various consultants and DHS.

Section 3 presents the results of the recent soil sampling investigation conducted by J & A.

Section 4 includes details on groundwater monitoring well drilling, construction, development, and groundwater sampling and analysis.

Section 5 is a general discussion of the chemistry, toxicology, and migration pathways of mercury and mercury fulminate.

Section 6 provides a summary of activities and presents results and conclusions.

- Actual or potential exposure to nearby human populations, animals, or the food chain from hazardous substances or pollutants or contaminants;
- Actual or potential contamination of drinking water supplies or sensitive ecosystems;
- High levels of hazardous substances or pollutants or contaminants in soils largely at or near the surface, that may migrate;
- Weather conditions that may cause hazardous substances or pollutants or contaminants to migrate or be released;
- Threat of fire or explosion;
- The availability of other appropriate federal or state response mechanisms to respond to the release; and
- Other situations or factors that may pose threats to public health or welfare or the environment.

The Phase II (PHEE) study will evaluate the available data in relation to these factors.

Phase III - If the Phase II evaluation indicates that removal action is appropriate, an Engineering Evaluation/Cost Analysis (EE/CA) or its equivalent will be performed. If collection of additional environmental samples becomes necessary, then a sampling and analysis plan will be prepared and submitted for regulatory review.

Under the NCP, possible Removal Actions may include:

- Fences, warning signs, or other security or site control precautions -- where humans or animals have access to the release;
- Drainage controls;
- Stabilization of berms and dikes -- where needed to maintain the integrity of the structure:
- Capping of contaminated soils or sludges -- where needed to reduce migration of hazardous substances or pollutants or contaminants into soil, ground or surface water, or air:
- Using chemicals and other materials to retard the spread of the release or to
  mitigate its effects -- where the use of such chemicals will reduce the spread of
  the release;
- Excavation, consolidation, or removal of highly contaminated soils from drainage or other areas -- where such actions will reduce the spread of, or direct contact with, the contamination; and

The references section cites documents consulted during the preparation of this report.

#### 1.1 SITE LOCATION

RFS is located at 1301 South 46th Street in Richmond, California 94804. Figure 1-1 is a general map of the RFS property and adjacent areas. As identified in the figure, the 150-acre property is located between Highway 580 and the Richmond Inner Harbor. Figure 1-2 is a site map of buildings and roads in the southern section of RFS, where the former mercury fulminate facility was also located. Figure 1-3 is a composite of current building locations, (shown in black in the figure) and former structures, (shown as a hatched line in the figure). The former mercury fulminate area appears as a circle. The approximate locations of the former buildings were determined from aerial photographs provided by RFS.

#### 1.2 SITE STATUS

The mercury fulminate facility was owned and operated by the California Cap Company during the 1920s. In 1981, RFS was included on the DHS list of abandoned hazardous waste sites because of the California Cap Company's explosives manufacturing activities. In addition, the Department of Fish and Game (DFG), San Pablo Sanitary District, and California Occupational Safety and Health Association (Cal OSHA) are aware that mercury may be present on the property.

To evaluate whether mercury was present at the site, DHS began an investigation of the RFS property in the general area of the mercury fulminate facility in 1981, at which time samples from five locations were collected and analyzed for mercury. Mercury was detected in two of these samples. In 1982, DHS collected 17 additional soil samples from the RFS facility and analyzed the samples for a large suite of inorganic constituents. Mercury was not detected in any of the samples collected by DHS in 1982. Two of these samples were from the same locations as the earlier DHS samples collected in 1981. In the 1982 sampling effort, none of the inorganic constituents in soil samples exceeded the Total Threshold Limit Concentrations (TTLC) set by DHS to identify a material as hazardous.

DHS concluded that no mercury remained on the RFS property and recommended that RFS should be removed from the list of abandoned hazardous waste sites. However, DHS continues to consider the facility as a "medium" priority until sampling results in areas previously tested can be verified.






FIGURE 1-1

REGIONAL MAP





Currently, RFS is not under any regulatory order from state or federal agencies to conduct any additional investigations of on-site mercury. However, because RFS is considering an addition of a new building near the former mercury fulminate facility, investigation of mercury contamination was initiated.

#### 1.3 CHRONOLOGY OF EVENTS

This section summarizes the history of pertinent activities occurring in the general area of the mercury fulminate facility from 1840 to 1990, including any related agency actions:

1840	Several explosives manufacturing companies were operating in the southern portion of current RFS property.
1900s	The California Cap Company bought a portion of the RFS property from the Hercules Powder Company.
1920	By 1920, California Cap Company had purchased all of the other explosives companies in the area, thereby becoming the only explosives manufacturer on the property.
1945	The California Cap Company ceased operations at the end of World War II.
1950	UC purchased the property. Under the terms of the sale, the California Cap Company was required to remove all hazardous material before UC would accept title. The company reportedly complied with this requirement, but the exact activities performed by California Cap Company are currently not known.
1981	In 1981, RFS was included on the DHS list of abandoned hazardous waste sites list because of the California Cap Company's explosives manufacturing activities. DFG, the San Pablo Sanitary District, and Cal OSHA agreed that fulminate of mercury might be present on the property, thereby presenting a risk of explosion and soil contamination. DHS began investigation of the RFS property in 1981.
<b>2</b> /18/81	In 1981, soil samples were collected from five locations by DHS. However, sampling and analytical methodologies performed are not known. Available data showed that mercury was found in two of these samples:

Sample 2, near the former Shell Manufacturing Area (see Figure 1-3), contained 105 parts per million (ppm) of mercury, and Sample 6, taken from the marsh, contained 23 ppm of mercury. A DHS report (DHS, 1982) indicated that arsenic, copper, zinc, and lead were also present in the RFS soil samples. However, the report did not identify concentrations.

- In 1982, DHS collected 17 soil samples from RFS and analyzed them for 29 inorganic constituents. Mercury was not detected in any of the samples. None of the other inorganic constituents analyzed for exceeded TTLCs set by DHS to identify a material as hazardous. At least two samples were analyzed for DDT; none was detected. DHS concluded that no mercury remained on the RFS property and recommended that RFS be removed from the list of abandoned hazardous waste sites.
- O3/10/83 The Regional Water Quality Control Board (RWQCB) wrote to UC stating that RWQCB had received copies of the DHS sampling results and considered the elevated levels of various metals (excluding mercury) to pose a threat to water quality.
- 02/23/84 RWQCB requested that UC submit a proposal for further study of potential water quality effects from on-site soil contamination.
- UC retained EAL Corporation (EAL) to analyze groundwater from two existing wells, and surface water, collected from the slough and marsh areas. The wells had been drilled previously for a research project and were completed approximately 90 feet below the ground surface.
- 04/05/84 EAL collected water samples in slough and marsh areas and sampled the two existing wells.
- 05/07/84 Report from EAL summarizing the April 5, 1984 sampling results.
- 07/05/84 RWQCB wrote a letter to UC requesting a water quality plan.
- 09/18/84 UC wrote a letter to RWQCB stating that, based on the previous sampling result and on surface runoff and infiltration information, RFS was not adversely affecting groundwater.
- 11/06/87 DHS requested copies of the existing sampling data.
- 11/17/87 UC sent requested data to DHS and asked that the site be removed from any hazardous site list.
- In 1987, DHS conducted another investigation of the facility, under U.S. Environmental Protection Agency (EPA) funding. The report concluded that no further EPA action or site investigation was necessary. DHS is continuing to consider the facility as a "medium" priority,

until sampling results in areas previously tested can be verified. DHS has indicated concern about the adequacy and quality assurance/quality control of its own sampling and analysis efforts, particularly the analytical results for soils sampled in 1982.

03/07/88

Aqua Science Engineers installed a groundwater monitoring well at RFS. Groundwater samples were collected to test for the possible presence of non-halogenated hydrocarbons that may have migrated from an existing underground fuel storage tank located outside of the study area.

06/03/88

Aqua Science Engineers submitted a report summarizing their results. Chemical analyses indicated that non-halogenated hydrocarbon concentrations in the groundwater ranged from below instrument detection levels for benzene, ethylbenzene, and total petroleum hydrocarbons (TPH), to 0.6 parts per billion (ppb) for xylene and 6.6 ppm for toluene.

10/18/88-11/02/88 CH2M Hill conducted additional sampling at RFS. The investigation was not required by DHS or RWQCB, but was initiated by UC to follow-up on previous sampling activities and to supplement and/or clarify analytical results.

During this investigation six soil sampling areas were identified and sampled throughout the RFS facility. Three of these sites were in the area occupied by the former mercury fulminate facility buildings. All samples collected were analyzed for metals. In addition, five samples (including the three samples from the former facility building sites) were analyzed for pesticides and PCBs. Levels of mercury above the TTLC were detected in the area of the former mercury fulminate facility. In addition to mercury, copper and zinc were found in these soils, but at concentrations below the TTLCs for these metals. Pesticides and PCBs were not detected in any of the soil samples analyzed.

12/89 to 3/90

J & A initiated a Phase I soil and groundwater sampling investigation in the general area of the former mercury fulminate facility as Phase I of the mercury investigation.

#### 2.0 INITIAL SITE EVALUATION

The following section concerns the previous work performed in the former mercury fulminate facility and Shell Manufacturing Area (see Figure 1-3).

#### 2.1 SITE DESCRIPTION

#### 2.1.1 Property History and Adjacent Land Use

#### 2.1.1.1 Property History

Beginning in the 1840s, several explosives manufacturing companies were operating in an area that is now the southeastern portion of RFS. At the beginning of the 1900s, the California Cap Company bought land and an explosive manufacturing operation from the Hercules Powder Company. By 1920, the California Cap Company had purchased all of the other explosives manufacturing companies in the area, thereby becoming the only explosives manufacturer on the property.

The California Cap Company ceased operations at the end of World War II in 1945. UC purchased the property in 1950. Under the terms of the sale, the California Cap Company was required to remove all hazardous materials before UC would accept the title. It is currently unknown if any corrective actions took place at that time.


RFS is currently owned and operated by UC. Throughout the years, several different buildings, including research laboratories and administrative offices, have been constructed on the property. Currently, RFS is considering an addition of a new building on the property. Figure 2-1 presents a possible future location for this proposed building. Due to the proximity of this proposed building to the former mercury fulminate facility, RFS decided to initiate an investigation of mercury at the former mercury fulminate facility.

#### 2.1.1.2 Adjacent Land Use

Current land uses adjacent to RFS are primarily industrial. A Safeway Distribution Center lies to the north, the Price Club and Bio Rad Labs are located to the east, and ICI Americas (ICI) lies to the west of RFS. Prior to 1987, the ICI Americas property was owned by Stauffer Chemical Company. The Richmond Inner Harbor lies to the south of the facility, along the shoreline of the City of Richmond.

Under regulatory direction from RWQCB, ICI completed a preliminary investigation of its property in December 1987. Under a Remedial Action Program designed and implemented by ICI, shallow groundwater onsite, found to be contaminated with pesticides, is currently being extracted and treated. The RWQCB has requested that ICI Americas perform a hydrogeologic investigation of the closed cinder landfill and associated ponds also formerly operated by Stauffer.

Recently the Richmond Inner Harbor has come under the scrutiny of EPA, DHS and the RWQCB, due to elevated concentrations of contaminants found in the water and biota of the harbor.



#### 2.1.2 Environmental Setting

#### 2.1.2.1 Regional Geology

RFS is located at the distal end of an alluvial plain that slopes in a southwesterly direction. To the northeast, towards the Berkeley Hills, the alluvial plain is transected by the Hayward Fault Zone (Jennings & Burnett, 1961). San Pablo Bay lies to the south of RFS.

The alluvial plain, upon which RFS is situated, is Quaternary in age representing relatively recent deposits. The lithology of the alluvial plain consists primarily of consolidated to unconsolidated clays, silts, sands, and gravels, and bordering Pablo Bay, organic-rich clays and silts. The total thickness of all these deposits ranges from surface deposits, where the alluvium thins against the Berkeley Hills, to approximately 300 feet in depth. These deposits are underlain by bedrock of the Mesozoic Franciscan Formation. The Franciscan Formation is a complex assemblage of serpentinite, greenstone, graywacke, chert, shale, sandstone, and schist, and is found on many ridges and mountains of the San Francisco Bay Region.

#### 2.1.2.2 Local Geology

Subsurface geology of the upper 100 feet in the area of RFS has been characterized by two boring logs, drilled for wells installed on RFS property (EAL, 1984). Both of these logs indicate that approximately 8 feet of clay directly underlie RFS. Interbeds of poorly sorted gravels, sand, and clay, extending to a depth of approximately 40 feet below ground surface lie beneath the clay layer. Clay to gravelly clay occurs from approximately 40 to 90 feet below ground surface. Sandy gravel was encountered from approximately 90 feet to the bottom of the borings at approximately 102 feet below ground surface. These lithologic descriptions are summarized in Table 2-1.

Table 2-1
Borehole Stratigraphy

Depth Below Surface	Lithology
(feet)	
0 - 8	Silty clay
8 - 40	Interbeds of poorly sorted gravels, sands, silts and clays
40 - 90	Clay to sandy gravelly clay
90 - 102	Sandy gravel

Four borings were drilled to a depth of 60 feet below ground surface, adjacent to the eastern boundary of RFS, on the ICI Americas property (ICI Americas, 1988). The logs from these borings were not available, but the local subsurface conditions were described as predominantly fine-grained sands, silts, and clays with minor amounts of gravel, found in poorly graded deposits of sand, silts, and clays (Hall, 1988). Discrete lithologies identified in these boreholes did not appear to be laterally or vertically extensive within the upper 60 feet.

#### 2.1.2.3 Surface Water

RFS is located adjacent to the Richmond Inner Harbor of San Pablo Bay. San Pablo Bay (in the northern portion of San Francisco Bay) forms the major surface water body to the south of the facility. Marshes border San Pablo Bay. Marshes at the southern extent of the facility drain south into San Pablo Bay through a single opening in a raised Santa Fe Railroad causeway.

No streams are present on the RFS property. Surface water runoff from the facility is collected and discharged through two storm drains, located at the southern extent of the property. These drains discharge into San Pablo Bay. To the west of RFS a tidal slough collects water from urban storm drainage throughout the City of Richmond. This slough is used by local industrial dischargers and is hydraulically connected with the marsh at the border of San Pablo Bay.

## 2.1.2.4 Regional Hydrogeology and Groundwater Quality

## 2.1.2.4.1 Regional Hydrogeology

RFS lies within the Alameda Bay Plain (ABP) Groundwater Basin (DWR, 1980). This groundwater basin extends southward from Richmond to Alvarado, between the Berkeley Hills and San Francisco Bay. The basin's hydrogeologic conditions are not well characterized because groundwater is currently not being used by local municipalities or for agricultural irrigation, due to its low permeability and the limited thickness of transmissive units.

Most groundwater recharge to the ABP groundwater basin in the Richmond area is probably from drainage through the beds of the Wildcat and San Pablo Creeks. Recharge from the infiltration of rainfall is believed to be limited due to the low permeability shallow clays and by the large amount of paved areas overlying the basin (Caltrans, 1978). Shallow groundwater probably eventually discharges into San Francisco Bay.

## 2.1.2.4.2 Local Hydrogeology

Local hydrogeology at the site is characterized from data obtained from monitoring wells installed on RFS property and the adjacent ICl Americas property (ICl Americas, 1988). Two groundwater monitoring wells are present at RFS. These wells are identified as MW-A and MW-1. MW-A is located approximately in the center of the RFS.

Monitoring well MW-1 is located behind Building 105, in the area of the former mercury fulminate facility (see Figure 1-2).

During the drilling of monitoring wells MW-A and MW-1, groundwater was first encountered at depths of 9.0 to 7.5 feet below ground surface, respectively. In the borehole for monitoring well MW-1, below a thick layer of clay, the first water-bearing unit composed of sand was located at a depth of 8 to 13 feet. This water-bearing unit was then screened. After monitoring well MW-1 was installed and allowed to equilibrate, the water level rose to 4.25 feet below the surface: indicating confined conditions. Because a thick clay layer exists below the surface, local groundwater probably occurs under confined to semiconfined conditions. Boring logs for MW-A and MW-1 are shown as Figures 4-2 and 4-3 in Section 4.0 of this report.

Geologic logs from boreholes drilled on RFS indicate that a second transmissive water-bearing unit of sandy gravel exists below a depth of 90 feet from the surface.

Groundwater in the vicinity of RFS is considered to flow from north-northeast to south-southwest, toward the Richmond Inner Harbor (Hall, 1988). Locally, a hydraulic cone of depression has been created by the groundwater extraction system currently operating at the adjacent ICI Americas facility. The center of this cone is located approximately 1,000 feet southeast of RFS border. Hall (1988) asserts that the actual cone of depression for the extraction system does not extend to RFS.

Several aquifer tests have been previously performed on RFS property. These tests were performed by the UC Sanitary Engineering Research Laboratory in 1954. A 3.5 foot gravelly aquifer was located at a depth of 90 feet below the surface. The average transmissivity was calculated from several tests to be 5,775 gal/day/ft. Dividing the calculated transmissivity by the measured thickness of the aquifer (3.5 feet), the estimated hydraulic conductivity, at 90 foot below the surface, is 1,760 gal/day/ft-squared (235 ft/day). These values indicate the gravelly aquifer located at a depth of 90 feet below the surface is a fairly transmissive zone. However, the shallower water-bearing zones are probably significantly less transmissive due to a greater presence of silts and clays in the aquifer material. Currently, no aquifer tests have been performed in the shallow water-bearing zone (8 to 13 feet below the surface) at RFS.

At the ICI Americas property, located adjacent to RFS, slug tests were conducted in the monitoring wells (Hall, 1988). These slug tests provided a rough estimate of hydraulic conductivity in shallow permeable zones of the upper 60 feet under the ICI Americas facility. The well completion details were not available for review. The calculated hydraulic conductivity values ranged between 0.4 to 17 gal/day/ft (0.05 to 2.3 ft/day). These values are well within the expected range for fine sand, silt, and clay (Freeze and Cherry, 1979).

The horizontal hydraulic gradient of the shallow groundwater at the ICI America property has been calculated to be approximately 0.004 (Hall, 1988). This was based upon a potentiometric surface map prepared from water level measurements from monitoring wells less than 60 feet deep. Using an estimated effective porosity of 0.25 and the hydraulic gradient of 0.004, shallow groundwater velocity is calculated to range between 0.3 to 13 feet/year.

surface. The results, presented in Table 2-3, indicate that most of the metal analytes in groundwater sampled exceeded EPA Sase Drinking Water Act standards.

# TABLE 2-3 Results of ICI Americas Well Sample Analyses (ICI Americas, 1987)

Constituent	Range in Concentration	Safe Drinking Water Act Water Standard
	(mg/L)	(mg/L)
Copper	<0.02 - 38	1.00°
Zinc	0.06 - 75	5.00 <sup>a</sup>
Arsenic	0.005 - 0.088	0.05
Iron	0.07 - 840	$0.30^{a}$
Cadmium	< 0.02 - 0.15	0.01
Lead	all $< 0.06$	0.05
Aluminum	<0.3 - 44	

a. Secondary Maximum Contaminant Level (aesthetic rather than health-based standard)

Limited water quality data for local groundwater encountered in borings less than 50-feet deep also exists in several Caltrans reports (1978 and 1979). Shallow groundwater from Caltrans borings taken in the vicinity of RFS was found to contain TDS ranging from 1,300 to 29,405 milligrams per liter (mg/l). In a more recent Caltrans investigation (1987), groundwater samples were collected from a series of 17 shallow wells to monitor the effects of a local freeway dewatering system. These wells were located to the north of RFS and further away from the influence of San Pablo Bay. With the exception of one well located near RFS which had TDS at greater than 1,000 mg/l, all other water samples contained TDS levels below 600 mg/l. In addition, the TDS concentrations in the deep wells that were constructed (to 50 feet) were generally less than the TDS concentrations in shallow wells (to 25 feet) which suggests that the shallower permeable zones are more prone to salt water intrusion from the Bay.

The RWQCB broadly defines a groundwater to have a beneficial use as a drinking water source when: 1) the groundwater is less than or equal to 3,000 ppm TDS; and 2) the aquifer produces a sufficient amount of water (approximately 150 gallons per day).

The groundwater sample collected from MW-1, in the area of the former mercury fulminate facility, showed no evidence of mercury contamination (detection limit of

0.001 mg/l). An additional analysis of a groundwater sample from monitoring well MW-1 (screened from 8 to 13 foot below surface), measured TDS at 1,300 mg/l.

#### 2.1.2.4.4 Surface Water Quality

As discussed earlier, surface water drainage from the facility is discharged into storm drains. These storm drains are eventually discharged into San Pablo Bay. Results of a previous analysis of samples from the east and west storm drains and from the sloughs are presented in Table 2-4.

TABLE 2-4

Results of Slough and Storm Drain Sample Analyses
(EAL Corporation, 1984)

	Sloughs (mg/L) Sample 3 Sample 1		RFS Storm Drains (mg/L) Sample 4 Sample 2		Surface Water Quality Criteria <sup>a</sup> (mg/L)
	South	North	East	West	
Mercury	<0.0005	0.0005	0.001	< 0.0005	0.0002
Copper	0.11	0.086	0.024	0.01	0.003
Zinc	0.072	1.1	0.56	0.76	0.095
Iron	2.4	0.55	0.11	0.54	•••
Lead	0.24	0.14	0.02	0.057	0.140

a. EPA National Ambient Water Quality Criteria to Protect Saltwater Aquatic Life, 1-hour average.

The results of the 1984 sampling by EAL indicate that with the exception of mercury, the concentrations of metals were higher in water from the sloughs than from the water in the storm drains that service RFS.

#### 2.1.2.4.5 Water Uses

In the area of South Richmond, Caltrans (1978) conducted a door-to-door survey for wells. The results of this survey identified over 100 wells in the general area of RFS. All of these wells are upgradient from RFS. Given the absence of historic agricultural or residential uses downgradient or proximate to RFS, it is not likely that unknown wells are present.

#### 2.2 SUMMARY OF THE PREVIOUS SOIL STUDIES (1981 - 1988)

#### 2.2.1 Previous Investigations

DHS conducted two rounds of sampling at RFS between 1981 and 1982. Sampling locations are presented on Figures 2-2 and 2-3.


In 1981, DHS collected soil samples from five locations. The results from the analyses of these samples identified mercury at two locations: Sample 2, near the former Shell Manufacturing Area, and Sample 6 collected in the marsh adjacent to San Pablo Bay. Results from analyzing Samples 2 and 6 for mercury identified 105 and 23 milligram/kilogram (mg/kg), respectively in these samples.


In 1982, DHS collected 17 additional soil samples from the RFS property and analyzed the samples for a large suite of inorganic constituents. Mercury was not detected in any of the samples collected by DHS. Two of these samples were from the same locations as the earlier Samples 2 and 6, collected in 1981 by DHS. None of the inorganic constituents analyzed exceeded TTLCs set by DHS to identify a material as hazardous. DHS concluded that no mercury remained on the RFS property and recommended that it should be removed from the list of abandoned hazardous waste sites.

In 1988, CH2M Hill collected additional samples at RFS. The purpose of this investigation was to follow-up on previous sampling activities and to supplement and/or clarify analytical results. Neither DHS nor RWQCB had requested that additional investigations be conducted at RFS. This program was an effort on the part of RFS to become more knowledgeable of previous hazardous waste management practices at the facility.

During this investigation three soil samples were collected in the general area of the former mercury fulminate facility. These samples are identified as SI-1, SI-2, and SI-3 on Figures 2-2 and 2-3. To collect these samples, test pits were excavated to approximately two feet. In each pit, samples were collected at various depths along the wall. These samples were then composited. After the sampling, the pit was refilled with the excavated soils. Prior to sample collection, the air in the pit was tested for the presence of volatile organic compounds using an HNu. No HNu measurements above background were recorded during sampling. All of the soil samples were analyzed for metals, pesticides, and PCBs. Pesticides and PCBs were not detected above analytical method detection limits. The concentrations of metals detected above analytical method detection limits are presented in Table 2-5.

Results from analysis of soil sample SI-1 indicate the presence of 260 mg/kg of mercury. Samples SI-2 and SI-3 also contained mercury, but at much lower concentrations of 0.53 and 0.34 mg/kg, respectively. All samples were collected from the site of the former mercury fulminate facility. The TTLC for mercury is 20 mg/kg, as specified in the California Code of Regulations, Title 26.





#### TABLE 2-5

## Results of 1988 Soil Samples Collected From the Former Mercury Fulminate Facility (CH2M Hill, 1988)

	SI-1	SI-2	SI-3	TTLC
	(mg/Kg)	(mg/Kg)	(mg/Kg)	(mg/Kg)
Mercury	260	0.53	0.34	20
Copper	<b>22</b> 3	47	16	2,500
Zinc	436	100	24	5,000

#### 3.0 CURRENT SOIL SAMPLING CONDUCTED BY JONAS & ASSOCIATES

#### 3.1 SAMPLING METHODS AND PROCEDURES


#### 3.1.1 Sample Collection and Analysis

From December 1989 through February 1990, four rounds of soil samplings were performed by J & A. Figure 3-1 shows sampling locations. The soil investigation was divided into three tasks. The purpose of task 1 was to define the horizontal extent of mercury in soil from 0 - 3 feet in depth range. Task 2 was conducted to determine concentration of mercury in surface soils (0 to 3 inches). Task 3 included soil sample collection from the borehole of the groundwater monitoring well, MW-1. Each task is described below.

## 3.1.1.1 Task 1 - Soil Sample Collection from 0 to 3 Feet Depth Range

In order to determine the horizontal extent of mercury in soil from 0 - 3 feet in depth range, 147 discrete soil samples (6 duplicates = 2 composites) were collected from 47 boreholes. At each borehole three discrete soil samples were collected; 0 - 1 foot, 1 - 2 feet, and 2 - 3 feet below the surface. During the on-site field effort, soil samples were obtained using a two-inch stainless steel hand auger. A sufficient amount of soil was collected for the selected laboratory analysis. The minimum sample volume required for laboratory analysis was 270 grams. Immediately after collection, samples were transferred to labeled, airtight sample containers. Sample containers from each borehole were sealed in an individual plastic bag. Samples were then placed into an insulated cooler for shipment to the laboratory, maintaining the temperature at approximately 4° Celsius. Upon receipt of samples by the laboratory, one composite sample was prepared per each sample location. A total of 49 composites were tested for total mercury (EPA Method 7471). Additional analyses are discussed below:

The composite soil sample from boring B-21 was analyzed for mercury using EP
Toxicity Extraction Method 3020/7470, to determine whether levels of mercury
in soil exceeded the designated federal hazardous waste level of 0.2 mg/l.



- The composite soil sample from boring B-18 was analyzed for soluble mercury using the Waste Extraction Test (WET), EPA Method 7471, in order to determine whether levels of mercury in soil exceeded the State hazardous waste level of 0.2 mg/l.
- Mercury fulminate is prepared by the reaction of mercury, ethyl alcohol and nitric acid. Therefore, composite soil samples from borings B-21, B37, B-39, and B-42 were analyzed for nitrate (EPA Method 300). These sample locations were selected because they are situated in the immediate area proposed for future redevelopment.
- Composite soil samples from borings B-21, B-22, B-23, and B-24 were analyzed for sulfate (EPA Method 300). In the past, the area sampled (except B-21) was reported to have been flooded by aqueous sulfate solutions from the ICI Americas' ponds. A composite sample from B-21 was analyzed for sulfate to obtain a background concentration.
- Composite soil samples from borings B-12, B-18, B-38, and B-39 were analyzed for copper and zinc (EPA Method 6010). These analyses were performed for confirmation of the previous sampling results obtained by CH2M Hill (Table 2-5). Similar to the CH2M Hill results, copper and zinc were detected below TTLC levels.

## 3.1.1.2 Task 2 - Surface Soil Sample Collection

In addition to the samples discussed above, ten surface soil samples were collected (0 - 3 inches in depth) and are identified as surface samples "SS". These samples were collected near existing boreholes identified in Figure 3-1. The surface samples collected and corresponding adjacent boreholes are identified in Table 3-1. Samples were collected using a stainless steel scooper, and were placed in jars inside an insulated cooler (maintaining them at approximately 4° Celsius), and shipped to the laboratory. All samples were analyzed for mercury (EPA Method 7471). Surface samples SS4 and SS9 (along with the composite samples collected from boreholes B-18 and B-39) were also analyzed for copper and zinc (EPA Method 6010).

3.1.1.3 Task 3 - Collection of Soil Samples from the Groundwater Monitoring Well (MW-1) Borehole

During drilling of the groundwater monitoring well MW-1, soil samples were collected from 2.5 feet to 11.5 feet and from 14 to 15 feet. Composite samples from 2.5 - 4.5 feet, 5 - 6 feet, 6.5 - 7.5 feet, 8 - 9 feet were analyzed for total mercury (EPA Method 7471). These samples were obtained in order to determine the vertical extent of mercury in soil. Soil samples were collected in 6 inch brass tubes and composited at the laboratory. Soil samples from 10.5 - 11.5 feet and 14 - 15 feet were not analyzed by the laboratory. They were collected as lithologic samples only.

Table 3-1
Surface Samples and Corresponding Borehole Locations

(J & A, 1990)

	Corresponding
Surface Samples	<b>Borehole</b>
	<b>Locations</b>
SS1	B-6
SS2	B-3
SS3	B-12
SS4	B-18
SS5	B-16
SS6	B-20
SS7	B-10
SS8	B-37
SS9	B-39
SS10	B-43

Field Chain-of-Custody records (Appendix A) completed at the time of sample collection accompanied the samples inside the cooler for shipment to the laboratory. The samples were properly documented on the field chain-of-custody record by the sampling team. Each cooler contained sufficient ice and/or ice packs to ensure that proper temperature was maintained and each was packed in a manner designed to prevent damage to sample containers. All coolers were promptly delivered for analysis to the Curtis & Tompkins, Ltd. analytical laboratories by the J & A field crew.

## 3.1.2 Decontamination and Post-Sampling Procedures

Decontamination of equipment occurred at a specific zone designated at the site. Sampling equipment was decontaminated prior to initial use, between sample locations, and at the completion of sampling activities. Items requiring decontamination were the stainless steel hand auger, a stainless steel spoon and a scooper, and the split spoon used during well drilling activity. A manual scrubbing to remove foreign material followed by a thorough cleaning was used for decontamination of the above items. All non-disposable equipment was decontaminated according to the procedures summarized below:

- o Manual scrub with non-phosphate soap solution followed by a tap water wash
- o Tap water rinse

- o Distilled/deionized water rinse
- o 10% nitric acid rinse (for metals only)
- o Distilled/deionized water rinse
- o Air dry
- o Distilled/deionized water rinse

#### 3.1.3 Sample Documentation

Sample documentation included field logbooks, sample labels, and Chain-of-Custody forms. All field documentation was written legibly in waterproof ink. Errors were crossed out with a single line, initialed, and dated.

#### 3.1.4 Sample Identification Numbers

Each sample was assigned a unique identification number that will allow retrieval of information regarding the sample. The sample identification number consists of three main parts, separated by a hyphen. The first part identifies the boring number, and is made up of the letter "B" and a one or two digit number representing the boring number. The second part represents the sampling depth. The third part, is the sampling date.

Example: B1-1-011690

Represents: Boring # 1, taken at the depth of 1 foot on January 16, 1990.

#### 3.1.5 Field Logbooks

A project field logbook was used to document the following:

- o Date and time of log entries;
- o Field conditions (weather, terrain, hazards, etc.);
- o Personnel present during field operations;
- o Decontamination procedures;
- o Waste disposal procedures, and a daily inventory of wastes present onsite;
- o Field measurements taken, instrumentation used, and frequency of instrument calibration;
- o Maintenance of instruments;
- o Information recorded on sample labels, as well as the site identification number and the sampling depth;
- o Any unusual sample characterization; and
- o Other specific considerations pertaining to sample acquisition.

#### 3.1.6 Sample Labels

A pre-printed adhesive label was affixed to each sample container (Figure 3-2). The information below was written on every sample label:

- o Project number;
- o Sample identification number;
- o Date of sampling;
- o Name of sample collector; and
- o Type of analysis requested.

Figure 3-2 Sample Label

Curtis & Topkins, Ltd., 2323 Fifth Street Berkeley, California 94710		
SAMPLE NO CUSTOMER SIGNATURE TYPE OF ANALYSIS	DATE	

## 3.1.7 Chain-of-Custody Records

A Chain-of-Custody record accompanied samples when they were shipped to the laboratory (Appendix A). The Chain-of-Custody documents the transfer of samples from one party to another. Additional information noted on this form includes:

- o Project number;
- o Sample identification number;
- o Date and time of sampling;

- o Type of sample; and
- o Type of analysis to be performed.

#### 3.1.8 Quality Control Samples

During Phase I of the investigation, two duplicate soil samples were collected and analyzed for mercury. The locations of the duplicate soil samples are presented in Figure 3-1 and identified as B-21 and B-45. Duplication of samples is a means of checking field and laboratory procedures. They are collected, numbered, and sealed in the same manner as the other samples.

#### 3.1.9 Sample Shipment

The J & A Technical Manager notified the Sales Manager at Curtis & Tompkins, Ltd. (CTL) a week before each round of sampling was scheduled to begin, so that the laboratory could prepare and ship the necessary coolers and sample bottles to the field team in advance. Each shipment of sampling supplies from CTL was accompanied by a cooler packing slip, which documented the number of coolers and the number and ... type of sample containers sent.

Samples were packaged for shipment in a cooler chilled with bags of ice. Sample jars were place in sealed Zip-lock bags. When possession of the samples was transferred, the individuals relinquishing and accepting custody wrote their names, the names of their organizations, and the time of custody transfer on the Chain-of-Custody Record(s).

### 3.2 DISCUSSION OF RESULTS OF SOIL ANALYSES

# 3.2.1 Results of Analyses for Mercury in Soil Samples

The results of the December 1989 and January 1990 J & A sampling efforts are presented in Figures 3-3 through 3-9. Figures 3-3, 3-4, and 3-5 present mercury concentrations from composite samples collected from the upper 3 feet of soil. Figure 3-6 presents mercury concentrations in the surface soil (0 - 3 inches). Figure 3-7 denotes the sulfate and nitrate concentrations in soil at 0 - 3 foot depth ranges. Figure 3-8 and 3-9 identify the concentrations of copper and zinc detected in soil samples collected from 0 - 3 foot depth ranges and from surface soil.

Results from 0 to 3 feet composite soil samples are presented in Table 3-2 and Figure 3-3.


Table 3 - 2


MERCURY IN 0 - 3 FEET OF SOIL
RICHMOND FIELD STATION
(J & A, 1990)


COMPOSITE SAMPLE <sup>a</sup>	MERCURY mg/kg	COMPOSITE SAMPLE <sup>a</sup>	MERCURY mg/kg
Bl	3.5	B24	41.3
B2	9.6	B25	0.46
B3	3.3	<b>B26</b>	0.80
B4	4.8	B27	1.2
<b>B</b> 5	19	<b>B28</b>	0.29
B6	22	<b>B29</b>	0.10
B7	<b>2</b> 6	B30	4.4
B8	28	B31	6.7
B9	61	B32	1.4
B10	90	<b>B33</b>	0.41
B11	<b>25</b>	B34	7.0
B12	180	<b>B</b> 35	3.0
B13	7.4	<b>B</b> 36	0.23
B14	8.1	B37	0.73
B15	27	<b>B38</b>	2.5
B16Db	<b>2</b> 8	B39D <sup>b</sup>	6.7
B17	140	B40	2.7
B18	630	B41	0.34
B19	27	B42	0.16
B20	20	B43	1.6
B21D <sup>b</sup>	44	B44	1.0
B22	11	B45Db	6.1
B23	2.2	B46	14
B24	41.3	B47	4.7
	****	B48	0.41


a. Composite samples identified by borehole number.


b. D = Duplicate

















Table 3 - 3

### MERCURY IN SURFACE SOIL RICHMOND FIELD STATION (J & A, 1990)

SAMPLE (BOREHOLE	MERCURY ) mg/kg	SAMPLE (BOREHOLE)	MERCURY mg/kg
SS2(B3)	4.7	SS1(B6)	3.9
SS7(B10)	21	SS3(B12)	90
SS5(B16)	24	SS4(B18)	<b>2</b> 40
SS6(B20)	1.2	SS8(B37)	13
SS9(B39)	1.1	SS10(B43)	0.92
9			

As seen in Figure 3-3, mercury concentrations ranged from 0.1 mg/kg to 630 mg/kg. The highest concentrations were located in the general area of Building 125 and Building 110. The lowest concentrations were in the area of the south-southwestern fence line and at Building 128. Three samples collected along this fence line (B-22, B-23 and B-24) ranged from 0.10 mg/kg to 0.46 mg/kg, with an average concentration of 0.26 mg/kg mercury. The seven borehole samples collected around Building 128 (B-38, B-39, B-41 through B-45) ranged from 0.16 mg/kg to 6.7 mg/kg, with an average concentration of 1.86 mg/kg mercury. Six samples (B-24, B-28, B-32, B-39 and B-44) defined the northern extent of sampling for mercury in soil. Results from these samples ranged from 0.29 mg/kg to 6.7 mg/kg.

Figure 3-4 presents the soil mercury results superimposed on the former mercury fulminate production area. As seen in this figure, high concentrations of mercury are located in this previous production area. Outside of this mercury production area, twenty borehole composite samples ranged in concentrations from 0.1 mg/kg to 6.7 mg/kg, with an average concentration of 2.3 mg/kg mercury. From this analysis there appears to be a very strong correlation between previous activities in the mercury fulminate production area and concentrations of mercury in shallow soil. Figure 3-5 shows that the main body of mercury in soil is localized in the area of the former mercury fulminate facility.

Duplicate samples were collected and analyzed. Samples B-16 and B-21 were collected from the same borehole. Mercury in these samples was, 28 mg/kg and 44 mg/kg, respectively. Samples B-39 and B-45 were also collected from the same borehole. Mercury in these samples was 6.7 mg/kg and 6.1 mg/kg, respectively. The differences in results between duplicate samples is probably due to a natural variation of mercury adsorption in the soil.

٠.

Results from sampling of the surface soil (0 - 3 inches) are presented in Table 3-3 and on Figure 3-6. As presented in Figure 3-6, mercury concentrations above 20 mg/kg are fairly localized in the area of the former mercury fulminate production facility.

During drilling of the monitoring well MW-1, 18 soil samples were collected from 2.5 to 11.5 feet and 14 to 15 feet depths. Four composite samples (2.5 - 9 feet) were analyzed for mercury. Mercury was detected in all four samples at a concentration range of 2.7 to 36 mg/kg. Discrete soil samples from 10.5 to 11.5 feet and 14 to 15 feet were collected for lithologic description only. These samples were not submitted to the laboratory for analysis.

# 3.2.2 Results of Analyses for Sulfate and Nitrate in Soil Samples

As Table 3-4 indicates, four composite soil samples (0 - 3 feet) were analyzed for sulfate and nitrate. The sample locations are shown in Figure 3-7. Concentrations of sulfate (SO<sub>4</sub><sup>-2</sup>) sampled east of Egret Way ranged from 170 mg/kg to 300 mg/kg and were collected from three boreholes. Another sample, collected behind Building 125, contained 5.9 mg/kg sulfate.

Table 3 - 4

SULFATE & NITRATE IN 0 - 3 FEET OF SOIL

RICHMOND FIELD STATION

(J & A, 1990)

COMPOSIT	TE SULFATE	COMPOSIT	E NITRATE
SAMPLE <sup>a</sup>	mg/kg	SAMPLE	mg/kg
B21	5.9	B21	44
B22	170	B37	18
B23	230	B39	13
B24	300	B42	11

a. Composite samples are identified by borehole numbers.

Three samples collected near Building 128 and one within the location of the former mercury fulminate facility were analyzed for nitrate  $(NO_3)$ . Nitrate in the samples collected near building 128 ranged in concentration from 11 to 18 mg/kg (Figure 3-7). The sample collected within the former mercury fulminate facility contained 44 mg/kg of nitrate. Currently, it is unknown whether the nitrate found represents a free radical of nitroglycerin  $(C_3H_5(ONO_2)_3)$ , mercury fulminate  $(C_2N_2O_2Hg)$ , or is from another source (garden fertilizer or previous sewage activity).

# 3.2.3 Results of Analyses for Copper and Zinc in Soil Samples

Table 3-5 and Figures 3-8 and 3-9 present the sample locations and analytical results for copper and zinc concentrations in composite samples B-12, B-18, B-38 and B-39. In addition, two surface samples were collected adjacent to the sample locations for B-18 and B-39. All samples were below the TTLC level of 2,500 ppm for copper, and 5,000 ppm for zinc.

#### Table 3 - 5

# COPPER AND ZINC IN 0 - 3 FEET OF SOIL RICHMOND FIELD STATION (J & A, 1990)

COMPOSITE	COPPER (TTLC:2,500) mg/kg	COMPOSITE	ZINC (TTLC:5,000)
SAMPLE		SAMPLE	mg/kg
B12	53	B12	65
B18	57	B18	640
B38	200	B38	170
B39	160	B39	180

# COPPER AND ZINC IN SURFACE SOIL RICHMOND FIELD STATION (J & A, 1990)

SAMPLE	COPPER (TTLC:2,500)	SAMPLE ZINC (TTLC:5,000)
(BOREHOLE)	mg/kg	(BOREHOLE) mg/kg
SS4(B18)	140	SS4(B18) 270
SS9(B39)	110	SS9(B39) 150

# 3.2.4 Results of EP Toxicity and Waste Extraction (WET) Tests for Mercury in Soil Samples.

In order to determine whether levels of mercury in soil exceeded the designated federal hazardous waste level (0.2 mg/l), an EP Toxicity Extraction (EPA 3020/7470) was preformed on the composite soil sample, B-21. The sample had originally shown 44 mg/kg of total mercury in soil. Results from EP Toxicity Extraction indicated that mercury in the extracted sample was not present above the detection limit of 0.01 mg/l.

In addition to this analysis, the State's Waste Extraction Test (Wet) was performed on the composite sample from borehole B-18 in order to determine whether levels of mercury in soil exceeded the State-designated hazardous waste level (0.2 mg/l). A mercury concentration of 1.3 mg/l was detected in the extracted waste. The results of the WET test indicate that mercury is present in these soils could potentially (under acidic conditions) leach out of soil at concentrations equal to or greater than 0.2 mg/l.

# 4.0 GROUNDWATER MONITORING WELL DRILLING, CONSTRUCTION, DEVELOPMENT, SAMPLING, AND ANALYSIS

To characterize the vertical stratigraphy and water table and to determine the presence or absence of mercury in groundwater at RFS, an additional groundwater monitoring well (MW-1) was installed downgradient from the former mercury fulminate facility by Aqua Science Engineers Inc. Figure 4-1 presents the well construction details of the monitoring well, MW-1. Figure 4-2 identifies the general lithologies encountered at the location of MW-1 and the results from the soil sample analysis.

A previous groundwater monitoring well was installed by Aqua Science Engineers on March 7, 1988 as part of an underground fuel storage tank investigation. Groundwater sample collected from this well showed no contaminant of concern. The location of this well is shown in Figure 2-2. This well is located upgradient from the former mercury fulminate facility. It will be considered an upgradient background well for this and all future investigations and will be referred to as MW-A. The well log description of MW-A is presented in Figure 4-3.

The methods and procedures for drilling, construction, development, and sampling of monitoring well MW-1 (installed pursuant to Phase I) are described in this section. Details regarding sample collection procedures, analysis, and analytical results are also included.

# 4.1 DRILLING METHODS

The soil boring for the monitoring well was drilled utilizing a hollow-stem auger. Hollow-stem auger drilling is accomplished through use of a hollow central shaft with an attached spiral scroll. Each section of auger is aligned so that a continuous scroll is formed. A bit is attached at the bottom of the first auger flight. Cuttings created by the bit are removed by the scroll as the auger stem is turned. This method is suitable for relatively shallow drilling in unconsolidated formations. A soil boring 13 feet deep was drilled with an auger rig utilizing an eight-inch outside diameter hollow-stem auger. This boring was completed as a monitoring well. The spoil was collected in drums and set aside for appropriate disposal.

Soil samples for lithologic description and chemical analysis were collected every foot, utilizing a split-barrel drive sampler. A total of 18 discrete soil samples were collected from 2.5 feet to 11.5 feet and from 14 to 15 feet. Soil samples were collected in brass tubes, labeled and placed in plastic bags. Samples were then placed into an insulated cooler for shipment to the laboratory, maintaining them at approximately 4° Celsius.


# JONAS AND ASSOCIATES INC.

WELL No.: MW1

PROJECT No.: UCRFS-101

Figure 4-1 MW-1 CONSTRUCTION DETAILS

PROJECT NAME: UNIVERSITY OF CALIFORNIA RICHMOND FIELD STATION BORING LOCATION: Behind Building 105 DATE: 2-25-90 BY: RJ



# JONAS AND ASSOCIATES INC.


Figure 4-2

WELL No .: MW1

PROJECT No.: UCRFS-101

# MW-1 STRATIGRAPHY AND SOIL ANALYTICAL RESULTS

PROJECT NAME: UNIVERSITY OF CALIFORNIA RICHMOND FIELD STATION BORING LOCATION: Behind Building 105 DATE: 2-25-90 BY: RJ






FIGURE 4-3

Upon receipt by the laboratory, one composite sample was prepared per depth ranges specified on the plastic bag. Four composite samples from 2.5 - 4.5 feet, 5 - 6 feet, 6.5 - 7.5 feet, 8 - 9 feet were analyzed for total mercury (EPA Method 7471). Details on sample collection and analysis are provided in Section 3 of this report.

#### 4.2 WELL CONSTRUCTION

The groundwater monitoring well MW-1 was completed with a 5-foot long PVC screen, with a 0.01-inch slot size. The screen was connected to a 10-foot long PVC riser pipe. All joints were flush threaded and no solvents or cements were used on the PVC. The pipe and screen were steam-cleaned before use. The screen penetrates the complete thickness of the aquifer.

A filter pack consisting of three-inch silica sand was placed in the annular space at the well screen and carried 2 feet above the top of the screen. A bentonite seal, one-half foot thick and composed of one-half inch pellets, was placed above the sand pack and the remaining annular space was filled with a neat cement-bentonite grout. Frequent measurements with a steel tape were made during placement of materials to ensure that proper amounts of material were placed and that seals were properly located to avoid cross-contamination. The grout used to finish the well consisted of neat Portland cement modified with sodium bentonite to reduce shrinkage. The ratio of cement to bentonite was approximately 20:1 on a weight basis. Water and then bentonite were added to the Portland cement.

The well head was completed approximately 1 foot above the grade. A three-foot protective metal casing that extends 1.7 feet below grade and 1.3 feet above grade was placed over the 1-foot riser. The metal casing includes a hinged steel lid with provision for a padlock.

When constructing the well, maximum effort was made to avoid contamination of the well construction materials. The PVC construction materials were procured clean from the fabricators. The following procedures were followed to prevent contamination:

- o All screen and casings were steam-cleaned before installation;
- o All filter-pack material was placed directly from the bag (spilled material was not to be taken from the ground and placed in the boring); and
- o The steel tape used to sound for depth during installation was steam-cleaned between borings.

Additional precautions taken during well construction included:

o Ensuring that no foreign material entered the well casing during construction;

- o Making frequent soundings when placing the filter pack, bentonite, and grout into the annular space;
- o Noting the total casing length in the borehole, the stickup, and the casing cutoff so that the total depth could be accurately calculated (i.e.: {(total casing length)-(stickup+cutoff)}=depth); and
- o Recording all final measurements, problems, and comments in the field log book.

## 4.3 WELL DEVELOPMENT

When well construction was complete and the grout column was cured for a minimum of 24 hours, well development was performed with a submersible pump. Well development was initiated by lowering a submersible pump into the well. The pump was connected directly to an outlet in one of the buildings nearest to the monitoring well. The pump was placed one foot above the bottom of the well. Approximately five well volumes were removed during development. Well volume was calculated using the . formula:

$$V = Pi(h/4) \{D_{c2}(1-n) + nD_{p2}\}$$

where V = volume of standing water in well, ft<sup>3</sup>

Pi = 3.14

 $D_p$  = diameter of filter pack, ft  $D_c$  = diameter of well casing, ft

n = porosity of the filter pack, decimal fraction

h = height of standing water in well, ft

The variable h was determined by subtracting the depth to water from the total well depth. The value n was 0.3. To convert the well volume to gallons, V was multiplied by 7.48. Water level and well depth were measured with an electric sounding device.

The monitoring well was pumped until the discharge was clear, colorless, and free of particulates. Water from equipment steam-cleaning and development were stored in 55-gallon drums.

# 4.4 SAMPLE COLLECTION PROCEDURES AND ANALYSIS

Upon the completion of well development, a teflon bailer was used to collect the sample. First, the teflon bailer was rinsed five-times with well water. Then the sample was directly poured from the bailer into a one-liter plastic sample bottle provided by Curtis & Tompkins laboratory. The bailer was decontaminated prior to and after sample collection. The sample bottle was sealed, labeled and kept at 4°C in an ice chest until transferred to the laboratory. The sample was filtered and preserved by the

laboratory upon receipt. The groundwater sample collected was analyzed for mercury (EPA Method 7470) and TDS (EPA Method 160.1).

# 4.5 RESULTS OF ANALYSES FOR MERCURY AND TOTAL DISSOLVED SOLIDS (TDS) IN WATER SAMPLE

Figure 4-4 presents the location of monitoring well MW-1 within the former mercury fulminate facility. MW-1 is located approximately 60 feet downgradient of the highest mercury concentration measured in soil (630 mg/kg in composite sample, B-18).

Mercury was not detected above the instrument detection limit of 0.001 mg/l in the groundwater sample. TDS was measured at 1,300 ppm in groundwater. Mercury was encountered in the soil to a depth of nine feet below the surface, as shown in Figure 4-2. The transmissive zone of silty sand, which was eventually screened for monitoring well MW-1, showed a mercury concentration in soil of 2.7 mg/kg. Concentrations above that zone were generally an order of magnitude higher.

The results of the groundwater sample analysis indicate that the mercury in the soil could be effectively bound and may not be releasing into groundwater. From a water quality perspective this is very encouraging, since mercury fulminate has been at this location for over forty years and it might be expected to be present in the groundwater. However, based on the groundwater analysis, no mercury was detected. This may be due to the insoluble property of mercury fulminate, as well as the presence of a thick clay layer (0 - 7.5 feet) that provides an organic environment for mercury to effectively bind with the organic fraction of the silts and clay. Additional groundwater samples will be collected as part of Phase II of the mercury investigation to augment the preliminary findings of Phase I.

# 5.0 THE CHEMISTRY, TOXICOLOGY, AND MIGRATION PATHWAYS OF MERCURY AND MERCURY FULMINATE


The discussion presented in this section is based on a general literature search. A public health and environmental risk assessment is proposed for phase II of the mercury investigation, at which time a thorough evaluation will be performed.

## 5.1 CHEMISTRY OF MERCURY AND MERCURY FULMINATE

# 5.1.1 Chemistry of Mercury

Mercury has an atomic weight of 200.59. At  $20^{\circ}\text{C}$ , its specific gravity is 13.546 and its vapor pressure is 0.0012 torr. The solubility of metallic mercury in pure water has been determined by Sanemasa (1975) to be 0.019 mg/l and 0.081 mg/l at 5°C and 30°C, respectively.

Under the usual conditions of temperature and pressure that occur in the environment, mercury can be present in any one of three different oxidation states. The most reduced form is the metal, which is a liquid at ordinary temperatures and which has a tendency to vaporize. The other two forms are the mercurous ion,  $Hg^{+1}$ , and the



mercuric ion Hg<sup>+2</sup>. Mercury can build many species. Some complex ions have considerable aqueous solubility, while others are relatively insoluble. Mercury also forms many stable organic complexes. Within a moderately oxidizing environment above pH 5, the predominant mercury species will be elemental mercury. Mildly reducing conditions can cause mercury to precipitate as a sulfate (cinnabar). Cinnabar has an extremely low aqueous solubility. In aquatic environments that are high in chloride, the solubility of mercury in oxygenated solutions may be greatly increased by the formation of mercuric chloride complexes (Garvis and Ferguson, 1972).

Equally important are the processes which produce the organic complexes of mercury. Two types of alkylated mercury compounds are formed in the environment:

- o Compounds with a single carbon-mercury bonds that act as substituted salts which are reasonably water-soluble.
  - An example is methyl mercuric chloride (CH<sub>3</sub>HgCl), which becomes CH<sub>3</sub>Hg<sup>+</sup> ion and Cl in solution.
- o Compounds with two carbon atoms attached to the mercury. These covalent mercury compounds are generally quite insoluble.
  - An example is dimethyl mercury (CH<sub>3</sub>HgCH<sub>3</sub>) which is volatile and is undisassociated in solution. Inorganic forms of mercury can be converted to organic forms by microbial action.

#### 5.1.2 Mercury Fulminate

Mercury fulminate  $\{Hg-(ONC)_2 \text{ or } C_2N_2O_2Hg\}$  has a molecular weight of 284.6 and a density of 4.42 g/cm<sup>3</sup>. Its percentage of nitrogen is 9.84%. It is a crystal at standard temperatures and pressure. Its heat of explosion is 355 Kilocalorie per kilogram (kcal/kg) or 1486 kilojoule per kilogram (kj/kg). Mercury fulminates deflagration point is 165°C or 330°F and its impact sensitivity is 1-2 newton meter (N m).

Mercury fulminate is practically insoluble in water. When dry, it is highly sensitive to shock, impact, and friction, and is easily detonated by sparks and flames.

Mercury fulminate is prepared by dissolving mercury in nitric acid, after which the solution is poured into 95% ethanol. After a short while, a vigorous gas evolution takes place and crystals are formed. When the reaction has ended, the crystals are filtered by suction and washed until neutral. The mercury fulminate product is obtained as small, brown-to grey pyramid-shaped crystals; the color is caused by the presence of colloidal mercury. Owing to its excellent priming power, its high brisance, and to the fact that it can easily be detonated, mercury fulminate was the initial explosive most frequently used prior to the appearance of lead azide. It was used in compressed form in the manufacture of blasting caps and percussion caps. Mercury fulminate is stored under water.

#### 5.2 FATE AND TRANSPORT OF MERCURY

Mercury and certain of its compounds can volatilize to the atmosphere from aquatic and terrestrial sources. In the atmosphere complex species of mercury commonly form. Precipitation is the primary mechanism for removal of mercury from the atmosphere. But photolysis can also breakdown airborne mercurials. Adsorption onto suspended and bottom sediments is commonly the fate of mercury in the aquatic environment. Sorption is strongest on organic materials. Mercury in soils generally complexes to organic compounds.

Virtually all mercury compounds can be remobilized by microbial conversion to methyl and dimethyl forms. Conditions reported to enhance biomethylation include large amounts of available mercury, large numbers of bacteria, the absence of strong complexing agents, near neutral pH, high temperatures, and moderately aerobic environments. Mercury is also strongly bioaccumulated by numerous mechanisms. Methylmercury is the most readily accumulated form of mercury in aquatic biota.

#### 5.2.1 Volatilization

Metallic mercury has a relatively high vapor pressure relative to other metals and commonly enters the atmosphere as several different gaseous compounds. The rate of vaporization of mercury and certain of its inorganic compounds decreases in the sequence  $Hg > Hg_2Cl_2 > HgCl_2 > HgS > HgO$ , according to Koksay and Bradshaw (1969). The microbial methylation of mercury also enhances evaporation of mercury.

### 5.2.2 Sorption

Mercury has an affinity for many surfaces. In water samples a major portion of the total mercury is found associated with particulate matter (Hinkle and Learned, 1969). Ramamoorthy and Rust (1976) found that sorption rates are highest in organic-rich soils, with sediment binding capacity closely related to organic content. They also found that desorption rates are low: less than one percent Hg leached from sediments after 70 hours of agitation in distilled water.

#### 5.2.3 Biotransformation

Mercury does take part in biological reactions which alter its mobility and toxicity. Bisogni and Lawrence (1975) evaluated the influences of inorganic mercury concentrations and speciation, pH, microbial activity, and redox potential on mercury methylation rates. They found that in a neutral pH the primary product of mercury methylation is methyl mercury. Methylation can occur under both aerobic and anaerobic conditions. More mercury methylation occurs when high number of bacteria are present. Therefore, highly organic sediments which favor bacterial growth have a higher methylation potential than inorganic sediments. Upon entering an aqueous system, mercurial compounds will tend to convert to methylmercury through microbial chemical transformation.

#### 5.3 TOXICOLOGY OF MERCURY

Mercury occurs as elemental mercury and as inorganic and organic compounds (mercury vapor, mercury liquid, mercury salts, short-chain alkylmercury compounds, alkoxyalkylmercury compounds and phenylmercury compounds), all having different toxicological properties.

Mercury is circulated naturally in the biosphere, 30,000 - 150,000 tons being released into the atmosphere by degassing from the earth's crust and the oceans. In addition, 20,000 tons of mercury are released into the environment each year by human activities such as combustion of fossil fuels and other industrial release. Yearly, 10,000 tons of mercury are produced for industrial use, a small part of which is used for synthesizing organic mercury compounds.

In nature, methylmercury is produced from inorganic mercury as a consequence of microbial activity. In fish, the major amount of mercury is methylmercury. Factors determining the methylmercury concentration in fish are: mercury content in water and bottom sediment; pH and redox potential of water; and species, age, and size of fish.

The toxic properties of mercury vapor are due to mercury accumulation in the brain causing neurological signs, involving an unspecific psychoasthenic and vegetative syndrome (micromercurialism). At high exposure levels mercurial tremor is seen, accompanied by severe behavioral and personality changes, increased excitability, loss of memory and insomnia. On a group basis, exposure levels are likely to be reflected in mercury concentrations in blood and urine. Occupational exposure to mercury concentrations in air above 0.1 milligram per cubic meter (mg/m³) may produce mercurialism. Micromercurialism has not been reported at concentrations below 0.01 mg/m³.

The acute and long-term action of mercuric salts, phenylmercury compounds and alkoxyalkylmercury compounds is likely to be gastrointestinal disturbance and renal damage - appearing as a tubular dysfunction with tubular necrosis in severe cases. The lethal dose in human is about 1 gram of mercuric salt. The mercury load on the kidney is best determined by analysis of renal biopsy. Mercury concentrations in the kidney between 10 and 70 mg/kg have been reported in poison cases involving renal injury. Levels below 3 mg/kg may be found in normal cases. Occasionally, mercuric compounds may cause idiosyncratic skin symptoms that may develop into severe exfoliative dermatitis or may cause glomerular nephritis. A specific form of idiosyncrasy, called acrodynia or pink disease, is seen in children. Most cases are associated with mercury exposure showing increased levels of mercury in urine.

The hazards involved in long-term intake of food containing methylmercury and in occupational exposure to methylmercury are due to the efficient absorption (90%) of methylmercury in man and the long retention-time (half-time 70 days) with accumulation of methylmercury in the brain. Chronic poisoning results in degeneration and atrophy of the sensory cerebral cortex, paresthesia, ataxia, hearing

and visual impairment. Prenatal exposure causes cerebral palsy and in less severe cases, psychomotor retardation. Methylmercury concentration in blood and hair reflects the body burden and the concentration in brain of methylmercury. Intake resulting in body burdens of less than 0.5 mg/kg body weight is not likely to give rise to detectable neurological signs in adults. This intake corresponds to blood values of less than 200 ug/l and mercury level in hair less than 50 mg/kg. However, this level of methylmercury exposure in pregnant females may result in inhibited brain development of fetus with psychomotor retardation of the child as a consequence. The highest level of methylmercury load in pregnant women, not associated with inhibition of fetal brain development, is not known.

The history of mercury toxicology has been reviewed by Goldwater (1964), the pharmacology and toxicology by Clarkson et al. (1972), the chemistry of mercury in biological systems by Carty and Malone (1979), the toxicology of methylmercury by a Swedish Expert Group (1971), and the toxicology and epidemiology by Friberg and Vostal (1972), by the Task Group on Metal Accumulation (1973), the Task Group on Metal Toxicity (1976), and by the WHO (1976,1980).

#### 6.0 SUMMARY AND RECOMMENDATIONS

#### 6.1 SUMMARY OF THE INVESTIGATION

UC has initiated a comprehensive investigation of potential mercury contamination at the former mercury fulminate facility at RFS in Richmond, California. Earlier investigations indicated the presence of mercury in soils. J & A was retained in December 1989, under order 0-200248-TR to begin Phase I of this investigation. The overall objectives of the entire investigation are to:

- Collect the necessary data to verify and characterize the presence of mercury contamination:
- Determine the extent of mercury contamination in soil;
- Evaluate groundwater quality within the study area;
- Evaluate the specific risks and hazards to public health and the environment that may result from the contamination; and
- Identify the appropriate cleanup criteria and provide a quantitative basis for selection of an effective removal action.

A summary of Phase I activities is presented below:

- o Collection of 147 discrete soil samples (0 3 feet) from 47 boreholes;
- o Analysis of 49 composite (0 3 feet) soil samples for mercury;
- o Collection and analysis of 10 surface soil samples (0 3 inches) for mercury;

- o Analysis of four composite soil samples for sulfate and nitrate;
- o Analysis of four composite (0 3 feet) and two surface soil samples (0 3 inches) for copper and zinc;
- o Analysis of one composite soil sample for mercury using EP Toxicity and WET extraction procedures;
- o Installation of one groundwater monitoring well, MW-1;
- o Collection of 18 discrete soil samples collected during the monitoring well drilling:
- o Mercury analysis of four composite soil samples collected from the monitoring well-borehole;
- o Collection and mercury analysis of one groundwater sample from monitoring well MW-1; and
- o Data analysis and report preparation.

#### 6.2 SUMMARY OF RESULTS

Current findings to date indicate that:

- o Mercury is present in the soil above designated state and federal hazardous waste levels;
- o Mercury was not identified in groundwater;
- o Copper and zinc are present in the soil below designated state and federal hazardous waste levels; and
- o Sulfate and nitrate were detected in soil in low concentrations.

#### 6.3 **RECOMMENDATIONS**

It is recommended that the following tasks be incorporated into Phase II of this investigation which is intended to serve as a preliminary Public Health and Environmental Evaluation (PHEE).

- Perform a preliminary Public Health and Environmental Evaluation (PHEE);
- Conduct field investigations as necessary to collect any additional data needed for the PHEE;
- · Continue groundwater sampling and analysis; and

• Using information developed by the PHEE provide a quantitative basis for selection of an appropriate removal action.

However, the exact scope of work for the Phase II of the investigation will be finalized after UC's meeting(s) with DHS and other regulatory agencies review of this report

#### REFERENCES

- Adrad, B., Project Officer, Department of Health Services. June 1988. Personal Communication.
- Aqua Science Engineers Inc. 1988. Soil Boring Investigation at 1301 South 46th Street, Richmond, California, June 3, 1988.
- California Regional Water Quality Control Board, Central Valley Region. 1986. The Designated Level Methodology for Waste Classification and Cleanup Level Determination.
- California Department of Health Services, Toxic Substances Control Division,
  Alternative Technology and Policy Development Section. 1986. The California Site
  Mitigation Decision Tree Manual. Sacramento, CA.
- California Department of Health Services. 1982. Hazardous Waste Surveillance and Compliance Report, Richmond Field Station, July 15, 1982.
- California Department of Health Services, Hazardous Materials Laboratory Section. 1989. Laboratory Report for Total Metal Analysis, RFS, Richmond, CA July, 28, 1989.
- California Regional Water Quality Control Board, Central Valley Region. 1988. A Compilation of Water Quality Goals, Prepared by John B. Marshack.
- CH2M Hill. 1987. Draft Workplan for the Phase I Remedial Investigation Feasibility Study, Summer Del Caribe, Richmond, California. Prepared for the California Department of Health Services, November 1987.
- Coate, A., CH2M Hill. 1988. Letter to Glaud Kuykendall, Richmond Field Station, summarizing results of environmental sampling at the University of California, Richmond Field Station, Dated December, 6, 1988.
- Cook, M., A. 1971. The Science of High Explosives, American Chemical Society Monograph Series, Robert E. Krieger Publishing Co., Inc., New York.
- Dodge, L.CH2M Hill. 1988. Letter to Glaud Kuykendall, Richmond Field Station recommending additional environmental investigations, Dated June 16, 1988.
- EAL Corporation. 1984. Analytical Results, RFS, May 7, 1984.
- Friberg, L. and et. al. 1986. Handbook on the Toxicology of Metals, Volume II: Specific Metals. Elsevier, New York.
- Gavis, J. and J.F. Ferguson. 1972. The Cycling of Mercury Through the Environment. Water Res. 6:989-1008.

- Gutstein, J., University of California, SEEHRL. 1989. Acidity in the Richmond Field Station Salt Marsh.
- Gutstein, J., University of California, Berkeley, SEEHRL. 1989. The Natural Areas of Richmond Field Station.
- Hall, L.E. 1988. Report of Hydrogeologic Investigation ICI Americas, Richmond, California, February 12, 1988. Richmond Field Station, University of California, Berkeley.
- Hall L.E., Environmental Services & Operations, ICI Americas Inc. Richmond, CA. 1989. Personal Communication, July.
- Helley, J. E. and K. R. LaJoie, U.S. Geological Survey. 1979. Flatland Deposits of the San Francisco Bay Region, California Their Geology and Engineering Properties, and Their Importance to Comprehensive Planning.
- Hinkle, M.E. and R.E. Learned. 1969. Determination of Mercury in Natural Waters by Collection on Silver Screens. pp.251-255. U.S. Geological Survey Professional Paper 650D, Washington, D.C. 299p.
- ICI Americas Inc. 1987. Assessment of Surface Impoundments at ICI Americas, Richmond, California, for TPCA, November 2, 1987.
- ICI Americas Inc. 1987. Report of Hydrogeological Investigation of Cinder Landfill at the Richmond Site. February 12, 1988.
- James, Roger, Regional Water Quality Control Board, San Francisco Bay Region. 1987. Letter to Mr. Joseph Odrzywolski, Stauffer Chemical Company, regarding the requirements of the Toxic Pits Cleanup Act, Dated <au 6. 1987.
- Jones & Stokes Associates, Inc. 1989. Wetland Delineations of Potential Sites for the EPA Laboratory, University of California, Richmond Field Station, Richmond, California, September 28, 1989.
- Kirk-Othmer. 1965. Encyclopedia of Chemical Technology, Volume 8, Second Edition, John Wiley & Sons, Inc.
- Kuykendall, Glaud. 1989 and 1990. Personal Communications.
- Meyer, R. 1981. Explosives. Second revised and extended edition, Verlag Chemie, Florida.
- Oddi, M. 1982. The South Richmond Marshes: An Ecological Analysis, in The East Bay Shoreline, Selected Environmental Issues: Chapter 7. Environmental Sciences Group, University of California, Berkeley.

- Ramamoorthy, S. and B.R. Rust. 1976. Mercury Sorption and Desorption Characteristics of Some Ottawa River Sediments. Can. J. Earth Sci. 13:530-536.
- San Francisco Bay Conservation and Development Commission. 1987. Staff Report on Water Quality in San Francisco Bay.
- Sanemasa, I. 1975. The Solubility of Elemental Mercury Vapor in Water. Bull. Chem. Soc. Japan. 48:1795-1798.
- Sax, I.N. 1989. Dangerous Properties of Industrial Materials, Van Nostrand Reinhold.
- Shively, J., College of Engineering. Richmond Field Station. 1982. Letter to Mr. Ben Gonzales, Environmental Health & Safety, regarding alleged hazardous chemical waste, Dated July 13, 1982.
- Singer, H. J., Regional Water Quality Control Board, San Francisco Bay Region. 1983. Letter to Frank Genta, Office Facilities Manager, Richmond Field Station, Dated March 10, 1983.
- Thomas, J.F., PhD, PE, Department of Civil Engineering, University of California, Berkeley. 1984. Letter to Lester Feldman, regarding removing Richmond Field Statrion from State Hazardous Waste Site List, Dated September 18, 1984.
- United States Department of Health and Human Services, Public Health Service.1987. Registry of Toxic Effects of Chemical Substances, 1985-86 Edition, Volume 3. Cincinnati, Ohio.
- University of California, Berkeley, Engineering Geosciences, Department of Materials Science and Mineral Engineering. 1986. Hydrogeologic Site Assessment of the Engineering Geoscience Well Field at the Richmond Field Station, Contra Costa County, California, December 4,1986.
- Ibid. 1987. Borehole Electromagnetic Survey of the Engineering Geoscience Well Field at Richmond Field Station.
- University of California, Richmond Field Station. Review of historical files and aerial photographs of Richmond Field Station, conducted by J & A in 1989.
- University of California, Sanitary Engineering Research Laboratory, Richmond Field Station. 1965. Final Report on Movement of Coliform Bacteria Through Porous Media, November 30, 1965.
- Ibid. 1954. Final Report in Laboratory Investigations of the Travel of pollution from Direct Recharge into Underground Formations, December 31, 1954.
- White, Charles, P.E., Dept. of Health Services. 1982. Letter to Mr. John Shively, College of Engineering, Richmond Field Station, regarding removal of the Richmond Field

- Station from the list of abandoned hazardous waste disposal sites, Dated December 17,1982.
- Woodward Clyde & Associates. 1980. Supplemental Geotechnical Exploration, Northern Regional Library Compact Shelving Facility, University of California, Richmond Field Station, December 16, 1980.
- Woodward, Clyde & Associates, 1953. Richmond Field Station Development, Step 1, Soil Investigation for Three Proposed Buildings, August 24, 1953.
- Woodward-Clyde-Sherard and Associates. 1962. Soil Investigation for the Proposed Towing Tank Extension, Building 275, Richmond Field Station, Richmond, California, January 25, 1962.
- Woodward-Clyde-Sherard and Associates. 1963. Soil Investigation for the Proposed Engineering Materials Laboratory Building, Richmond, California, January 25, 1963.

# APPENDIX A

Jonas & Associates Soil Sampling Laboratory Results

and

Chain-of-Custody Forms

JONAS & ASSOCIATES INC.

# APPENDIX A

Jonas & Associates Soil Sampling Laboratory Results

and

Chain-of-Custody Forms



# ■ Curtis & Tompkins, Ltd., Analytical Laboratories. Since 1878

2323 Fifth Street, Berkeley, CA 94710, Phone (415) 486-0900

DATE RECEIVED: 12/27/89 DATE REPORTED: 01/04/90

PAGE 1 OF 5

LAB NUMBER: 19050

CLIENT: JONAS & ASSOCIATES, INC.

REPORT ON: 21 SOIL COMPOSITES

PROJECT #: J&A-UCRFS-101 LOCATION: UCRFS-SOIL BORING

RESULTS: SEE ATTACHED

OA/OC Officer

Laboratory Director

Berkeley Wilmington Los Angeles



LABORATORY NUMBER: 19050

CLIENT: JONAS & ASSOCIATES, INC.

PROJECT #: J&A-UCRFS-101

LOCATION: UCRFS-SOIL BORING

DATE RECEIVED: 12/27/89

DATE ANALYZED: 01/02/90

DATE REPORTED: 01/04/90

PAGE 2 OF 5

ANALYSIS: MERCURY

PREPARATION METHOD: EPA 3020 METHOD REFERENCE: EPA 7471

LAB ID	SAMPLE ID	RESULT	UNITS	REPORTING LIMIT
19050 - 1,2,3	B11122689/ B12122689/	3.5	mg / Kg	1.0
19050- 4,5,6	B13122689 B21122689/ B22122689/	9.6	mg / Kg	1.0
19050- 7,8,9	B23122689 B31122689/ B32122689/	3.3	mg /Kg	0.10
19050- 10,11,12	B33122689 B41122689/ B42122689/	4.8	mg/Kg	0.10
19050- 13,14,15	B43122689 B51122689/ B52122689/	19	mg / Kg	10
19050-	B53122689 B61122689/	2 2	mg/Kg	10
16,17,18	B62122689/ B63122689			

QA/QC:

RPD. %	<1	
RECOVERY, %	99	



LABORATORY NUMBER: 19050

CLIENT: JONAS & ASSOCIATES, INC.

PROJECT #: J&A-UCRFS-101 LOCATION: UCRFS-SOIL BORING DATE RECEIVED: 12/27/89
DATE ANALYZED: 01/02/90
DATE REPORTED: 01/04/90

PAGE 3 OF 5

ANALYSIS: MERCURY

PREPARATION METHOD: EPA 3020 METHOD REFERENCE: EPA 7471

LAB ID	SAMPLE ID	RESULT	UNITS	REPORTING LIMIT
19050-	B71122689/	26	mg / Kg	10
19,20,21	B72122689/ B73122689			
19050-	B81122689/	28	mg/Kg	10
22,23,24	B82122689/ B83122689			
19050- 25,26,27	B91122689/ B92122689/	61	mg/Kg	10
20,20,27	B93122689			
19050- 28,29,30	B101122689/ B102122689/	90	mg/Kg	10
,,	B103122689			
19050- 31,32,33	B111122689/ B112122689/	25	mg/Kg	10
•	B113122689			
19050- 34,35,36	B121122689/ B122122689/	180	mg/Kg	10
	B123122689			

QA/QC:

RPD, %	<1	
RECOVERY, %	99	



LABORATORY NUMBER: 19050

CLIENT: JONAS & ASSOCIATES, INC.

PROJECT #: J&A-UCRFS-101

LOCATION: UCRFS-SOIL BORING

DATE RECEIVED: 12/27/89

**DATE ANALYZED: 01/02/90** DATE REPORTED: 01/04/90

PAGE 4 OF 5

ANALYSIS: MERCURY

PREPARATION METHOD: EPA 3020 METHOD REFERENCE: EPA 7471

LAB ID	SAMPLE ID	RESULT	UNITS	REPORTING LIMIT
19050-	B131122689/	7.4	mg / Kg	1.0
37,38,39	B132122689/ B133122689			
19050- 40,41,42	B141122689/ B142122689/	8.1	mg/Kg	5.0
40,41,42	B143122689			
19050- 43,44,45	B151122689/ B152122689/	27	mg/Kg	10
· • • •	B153122689			
19050- 46,47,48	B161122689/ B162122689/	28	mg/Kg	10
	B163122689			
19050- 49,50,51	B171122689/ B172122689/	140	mg / Kg	1 0
10070	B173122689	420	. ***	4.50
19050- 52,53,54	B181122689/ B182122689/ B183122689	630	mg/Kg	450

QA/QC:

RPD, %	<1
RECOVERY, %	99



LABORATORY NUMBER: 19050

CLIENT: JONAS & ASSOCIATES, INC.

PROJECT #: J&A-UCRFS-101 LOCATION: UCRFS-SOIL BORING DATE RECEIVED: 12/27/89


DATE ANALYZED: 01/02/90 DATE REPORTED: 01/04/90

PAGE 5 OF 5


ANALYSIS: MERCURY

PREPARATION METHOD: EPA 3020 METHOD REFERENCE: EPA 7471


LAB ID	SAMPLE ID	RESULT	UNITS	REPORTING LIMIT
19050- 55,56,57	B191122689/ B192122689/	27	mg / Kg	10
33,30,37	B1921220897 B193122689			
19050-	B201122689/	2 0	mg/Kg	10
58,59,60	B202122689/ B203122689			
19050-	B211122689/	44	mg/Kg	10
61,62,63	B212122689/		<b>3</b> . <b>3</b>	
	B213122689			




			Laboratory Notes :		2 7		, r		- ·		-	*	Wate Soil Wast Oil #Con H <sub>2</sub> SO HNO <sub>3</sub> Ice	Matrix Methox	Client Contact Romens	tion U	iith Street y, Californi 486-0900	Curtis & Tomp
					81212268989122611	11122689891	W 3 1 2 26 24 84 1 2 26 11 :	1122689 8912261	8:3122689891122611	12767989127611	111726897912261	0 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	Yr Mo Dy T		Recorder	5-501 b	Samplers	Tompkins, Ltd   chain of custody for
Helinquished by: (signature ) Date/Hr Dispatched by: (signature ) Date/Hr	Relinquished by: (signature ) Date/Hr	by: (sign	Chain of Ci	$\vdash$	:42 > abox-	: 45\ 30me 40	30 Summan	· 30 The Fullish Anglysis	אפזעות ח		Make in			SAMPLE NOTES 601/8010 602/8020	Roman Jonas		Ramene Jones 17	study form -
Received by (signature) Received for Lab by (signature)  (UNUMATION 1921/84 2000)	Received by (signature )	Received by (signature )	Chain of Custody Record				×			X			EPA EPA Titl EPA TPH Benz Oil	624/8240 625/8270 e 22 Met. PP Metal: Method- ene-Tolucand Greas 608/8080	s (# ) ene-Xyle se Pestici			ANALYSIS REQUESTED



Received for Lab by (signature) Received by (signature) Received by (signature) Received by (signature) Received by (signature) ANALYSIS REQUESTED Pesticides & PCB's 0808/809 **EPA** Grease and 770 Benzene-Toluene-Xylene(s) aventh Chain of Custody Record три мегрод-PP Metals Title 22 Metals 0758/829 EPA 078/729 Eby P8/12/21 asia! 0208/209 EPA Relinquished by: (signature ) Date/Hr Dispatched by: (signature ) Date/Hr EPA 0108/109 autore SAMPLE NOTES z Chain of Lustouy Form ş かった Romen Samplers Hankeya Recorder Renums Time ン Sampling Date δ £ ナロ:Joan Sample Number Bruco Curtis & Tompkins, Ltd 0 Job Description VCRFS-Soil Job Number 3+ A - UCRF 2 2 Client Contact Kontender Other Preserved Мопе Berkeley, California 94710 (415) 486-0900 Method aoI EQ7H DOS2 H Laboratory Notes 2323 Filth Street #Containers Matrix TTO Maste **TTOS** Mater





		23 Be	23 rke	Filt Filt ley, 48	ո Տ Ca	tree lifor	t nia			•	ki	n:	<u> </u>	L	t			<b>-</b> [	. In	18								dy Form	<b>-</b> [-			=	ANA		SIS		SUE:	PCB's	14/			
	Jol	b [ b N ent	um	be		\Z\	A	-	ΰc	C R	? E	s -	<u>- )</u>	0	1		· ( )	'n				R	lecc	orc	der,	R	57	nena Jona					5	~ *		ne-Xvlene(s)		sticides	MITHOR 74			
	Water	Waste	atri	П		die.		letho	Vec	Other	1	Sar	7	R.	5) hr	7		er			T		ling			e		SAMPLE NOTES		EPA 601/8010		A 624/8240	irle 22 Meta	EPA pp Metals (* )	PH Method-	enzene-Toluer	Oil and Grease	608/8080	METCULY - 1			
R3)	13	ň   <u>i</u> ≊ ∡/	Ö		_	¥ :=	:   3		Ł	P	上	_		7	3	) 	<b>—</b>	_	-	4	M	р 2	-	-,-		T	<del>-</del>		ŀ	ដីជ	1 6	EPA E	+	12		ă	Ö		₹	+	╀	•
ן נגייו	H	Ŧ	╁	H	+	1	$\dagger$	r	十	+	$\dagger$	卜	H	व	12	1	H	$\vdash$	٤	╫	-		14	7	7	$\dagger$	$\dagger$	some as page 3	ŀ	十	十	+	十	+	Н		H	$\vdash \uparrow$	+	+	+	
20 6	廿	止	上				1	$\perp$	土	土	32	3	$\overline{\Sigma}$	2	2	Z	8	9	8	19	_	Z	2		4 :	1	3 5					工	I	I					X	I		
61	Н.	1	1	$\sqcup$	4	_	_	<u> </u>  X	1	$\downarrow$	32	1	1	2	2	6	8	13	8	19	1	2.	ZK	4	3	4				4	4	4	$\downarrow$	$\downarrow$			Ш	Ц	]	$\bot$	$\perp$	
216	H	Ц-	╀	$\vdash$	-	+	╀	$-\frac{1}{\lambda}$	+	╀	1	<u>2</u>   3	H	-	F	F	┞	╁	┝	╀	-	F	H	+	+	1	4	Some au Page		╬	╀	+	╀	╁	₽	$\vdash$	$\vdash$	$\vdash \vdash$	4	+	╀	
• -	H	+	十	H	┪	+	╅	+	十	十	ť	尸		H		├-	┢	†-	┢	╁╴	十	4	Ħ	+	+	十	十		ŀ	┪	十	十	十	十	H	H	H	$\sqcap$	十	+	╁	,
									1	I	I												П	I		I	I					工	1	工			口	口	丁	工		
	H	_	╀	$\sqcup$	4		4	4	╀	+	1	Ļ	igspace	_	L	<u> </u>	L	╀	L	╀	_	L	$\sqcup$	4	_ _	4	4			4	4	4	1	1			Ш		4	4	1	
	H	+	╁	╂┨	$\dashv$	╁	╀	╁	╀	+	╁	╀	╀	-	╀	┞	┝	╀	╀	╀╌	╁╴	┢	╂┼	+	+	╁	╁		ŀ	+	╁	╁	╁	+	╁	┞┤	H	$\vdash$	+	+	╀	
	<u> </u>	•	<del>-</del>	1					<del></del>			_		_		<u> </u>	_					<u> </u>		_			ΙΓ	Chain					<u>-</u>	<u>_</u>							<u></u>	
	La	ıboı	rate	ory	No	otes	3 :	;																			-	Chain				100						1.		<del>_</del>		
																					•						11	Relinquished by: (signature ) t			92		н	ece	IVEG	ю	(5	igna	ature	<b>,</b> ,		
																												Romano Jonas 17, Relinquished by: (signature ) [		•	2/			ace!			le		ature			•
																											11	tomiquonos by, toignaturo y s			ŀ		•••	000.	,,,,	υ,	101	· gr · c	iio, e	•		
																												Relinquished by: (signature ) [	Date	e/Hr			R	ecel	ved	l by	(s	lgna	ature	, )		
	İ																										-	Relinquished by: (signature ) [	Date	e/Hr			R	ecel	ved	by	(s	igna	ature	<del>, )</del>		•
																											$\ \cdot\ _{L^{2}}$	Dispatched by: (signature ) Di	ate	/Hr	1	las	R	ecei	ved	19	Lei	b b)	(5	ilgna		



# Curtis & Tompkins, Ltd., Analytical Laboratories, Since 1878

2323 Fifth Street, Berkeley, CA 94710; Phone (415) 486-0900

DATE RECEIVED: 12/27/89 DATE REPORTED: 01/17/90

PAGE 1 OF 2

LAB NUMBER: 19172

CLIENT: JONAS & ASSOCIATES

REPORT ON: 1 SOIL COMPOSITE

PROJECT #: J&A-UCRFS-101 LOCATION: UCRFS-SOIL BORING

RESULTS: SEE ATTACHED

QA/QC Office

Laboratory Dir

Los Angeles



LABORATORY NUMBER: 19172
CLIENT: JONAS & ASSOCIATES
PROJECT #: J&A-UCRFS-101
LOCATION: UCRFS-SOIL BORING

DATE RECEIVED: 12/27/89
DATE REQUESTED: 01/12/90
DATE ANALYZED: 01/17/90
DATE REPORTED: 01/17/90

PAGE 2 OF 2

ANALYSIS: MERCURY

METHOD REFERENCE: EP TOXICITY EXTRACTION/EPA 3020/EPA 7470

\_\_\_\_\_\_\_

LAB ID SAMPLE ID RESULT UNITS DETECTION LIMIT

19172-1 B211 122689/ ND mg/L 0.01

B212 122689/ B213 122689

ND = NONE DETECTED

QA/QC:

RPD, % <1
RECOVERY, % 93



### Curtis & Tompkins, Ltd., Analytical Laboratories, Since 1878

2323 Fifth Street, Berkeley, CA 94710, Phone (415) 486-0900

DATE RECEIVED: 01/19/90 DATE REPORTED: 01/31/90

PAGE 1 OF 7

LAB NUMBER: 19371

CLIENT: JONAS & ASSOCIATES

REPORT ON: 24 SOIL COMPOSITES

PROJECT #: J&A-UCRFS-101

LOCATION: UCRFS-SOIL BORING PHASE II

RESULTS: SEE ATTACHED

Jahoratory Director

Berkeley Wilmington Los Angeles



LABORATORY NUMBER: 19371 CLIENT: JONAS & ASSOCIATES PROJECT #: J&A-UCRFS-101

LOCATION: UCRFS-SOIL BORING PHASE II

DATE RECEIVED: 01/19/90
DATE ANALYZED: 01/25/90
DATE REPORTED: 01/31/90

PAGE 2 OF 7

ANALYSIS: MERCURY

PREPARATION METHOD: EPA 3020 METHOD REFERENCE: EPA 7471

\_\_\_\_\_\_

LAB ID	SAMPLE ID	RESULT	UNITS	DETECTION LIMIT
	./			
19371-	B221011690/	11	mg / Kg	1.0
1,2,3	B222011690/			
	B223011690			
19371-	B231011790/	2.2	mg/Kg	1.0
4,5,6	B232011790/			
	B233011790			
19371-	B241011790/	1.3	mg/Kg	0.10
7,8,9	B242011790/		• •	
	B243011790			
19371-	B251011690/	0.46	mg/Kg	0.10
10,11,12	B252011690/			
	B253011690			
19371-	B261011790/	0.80	mg / Kg	0.10
13,14,15	B262011790/			
	B263011790			
19371-	B271011790/	1.2	mg/Kg	0.10
16,17,18	B272011790/			
	B273011790			
19371-	B281011790/	0.29	mg/Kg	0.10
19,20,21	B282011790/			
	B283011790			

### QA/QC:

	**=====================================
RPD, %	13
RECOVERY, %	99
	:======================================



LABORATORY NUMBER: 19371

CLIENT: JONAS & ASSOCIATES

PROJECT #: J&A-UCRFS-101

DATE RECEIVED: 01/19/90

DATE ANALYZED: 01/25/90

DATE REPORTED: 01/31/90

LOCATION: UCRFS-SOIL BORING PHASE II PAGE 3 OF 7

\_\_\_\_\_\_

ANALYSIS: MERCURY

PREPARATION METHOD: EPA 3020 METHOD REFERENCE: EPA 7471

LAB ID	SAMPLE ID	RESULT	UNITS	DETECTION LIMIT
19371-	B291011790/	0.10	mg / Kg	0.10
22,23,24	B292011790/			
	B293011790			
19371-	B331011790/	0.41	mg/Kg	0.10
25,26,27	B332011790/			
	B333011790			
19371-	B341011790/	7.0	mg / Kg	1.0
28,29,30	B342011790/			
	B343011790/			
19371-	B301011990/	4.4	mg / Kg	1.0
31,32,33	B302011990/			
	B303011990	•		
19371-	B311011990/	6.7	mg / Kg	1.0
34,35,36	B312011990/			
	B313011990			
19371-	B321011990/	1.4	mg / Kg	1.0
37,38,39	B322011990/			
	B323011990			
19371-	B371011990/	0.73	mg /Kg	0.10
40,41,42	B372011990/			
	B373011990			

#### OA / OC

=======================================	
RPD, %	13
RECOVERY, %	99



LABORATORY NUMBER: 19371

CLIENT: JONAS & ASSOCIATES

PROJECT #: J&A-UCRFS-101

DATE RECEIVED: 01/19/90

DATE ANALYZED: 01/25/90

DATE REPORTED: 01/31/90

LOCATION: UCRFS-SOIL BORING PHASE II PAGE 4 OF 7

\_\_\_\_\_\_\_

ANALYSIS: MERCURY

PREPARATION METHOD: EPA 3020 METHOD REFERENCE: EPA 7471

LAB ID	SAMPLE ID	RESULT	UNITS	DETECTION LIMIT
19371-	B381011990/	2.5	mg /Kg	1.0
43,44,45	B382011990/		<b>3</b> . <b>3</b>	
,,	B383011990			
19371-	B391011990/	6.7	mg/Kg	1.0
46,47,48	B392011990/			•
, ,	B393011990			
19371-	B451011990/	6.1	mg/Kg	1.0
49,50,51	B452011990/			
	B453011990			
19371 -	B351011990/	3.0	mg/Kg	1.0
52,53,54	B352011990/			
	B353011990			
19371-	B441011990/	1.0	mg/Kg	0.10
55,56,57	B442011990/		<del>-</del> -	
	B443011990			
		•		
19371 -	B431011990/	1.6	mg/Kg	0.10
58,59,60	B432011990/			
	B433011990			
19371 -	B421011990/	0.16	mg/Kg	0.10
61,62,63	B422011990/			
	B423011990			
19371 -	B411011990/	0.34	mg/Kg	0.10
64,65,66	B412011990/		_	
	B413011990			

***************************************	
RPD, %	13
RECOVERY %	9 9



LABORATORY NUMBER: 19371

CLIENT: JONAS & ASSOCIATES

PROJECT #: J&A-UCRFS-101

DATE RECEIVED: 01/19/90

DATE ANALYZED: 01/25/90

DATE REPORTED: 01/31/90

LOCATION: UCRFS-SOIL BORING PHASE II PAGE 5 OF 7

ANALYSIS: MERCURY

PREPARATION METHOD: EPA 3020 METHOD REFERENCE: EPA 7471

\_\_\_\_\_\_\_

LAB ID	SAMPLE ID	RESULT	UNITS	DETECTION LIMIT
19371- 67,68,69	B401011990/ B402011990/ B403011990	2.7	mg / Kg	1.0
19371- 70,71,72	B361011790/ B362011790/ B363011790	0.23	mg / Kg	0.10 .

RPD, %

RECOVERY, %



LABORATORY NUMBER: 19371

CLIENT: JONAS & ASSOCIATES

PROJECT #: J&A-UCRFS-101

DATE RECEIVED: 01/19/90

DATE ANALYZED: 01/31/90

DATE REPORTED: 01/31/90

LOCATION: UCRFS-SOIL BORING PHASE II PAGE 6 OF 7

\_\_\_\_\_

ANALYSIS: SULFATE

METHOD REFERENCE: EPA 300.0

LAB ID	SAMPLE ID	RESULT	UNITS	DETECTION LIMIT
19371- 1,2,3	B221011690/ B222011690/ B223011690	170	mg / Kg	7 0
19371- 4,5,6	B231011790/ B232011790/ B233011790	230	mg/Kg	7 0
19371- 7,8,9	B241011790/ B242011790/ B243011790	300	mg / Kg	7 0

QA	1	QC	;

RPD, %
RECOVERY, %

95



LABORATORY NUMBER: 19371

CLIENT: JONAS & ASSOCIATES

PROJECT #: J&A-UCRFS-101

DATE RECEIVED: 01/19/90

DATE ANALYZED: 01/31/90

DATE REPORTED: 01/31/90

LOCATION: UCRFS-SOIL BORING PHASE II PAGE 7 OF 7

ANALYSIS: NITRATE


METHOD REFERENCE: EPA 300.0


LAB ID	SAMPLE ID	RESULT	UNITS	DETECTION LIMIT
19371- 40,41,42	B371011990/ B372011990/ B373011990	18	mg / Kg	10
19371- 46,47,48	B391011990/ B392011990/ B393011990	13	mg / Kg	10
19371- 61,62,63	B421011990/ B422011990/ B423011990	11	mg / Kg	10


QA/QC:

RPD, % <1
RECOVERY, % 101


	dies beompkins, Llu		<u> </u>	. ANALYSIS NECUESTED
	2323 Filth Street Berkeley, California 94710 (415) 486-0900	Samplers <u>Ro</u>	mena James	3C3's
	Job Description CCRFS _ Soil	Olivisa Plane II		· I · I · I · I · I · I · I · I · I · I
	Job Description CCRFS _ So. Do Job Number 3+1-UCRFS _ 161			
	Client Contact Ramence Jonas	Recorder <u>Ro</u>	( )	als s (* ) ane-Xvlene(s) se ?esticides 6
	Matrix Method Sample Number	er Sampling Date	9010	8020 3240 3270 3270 3270 36- 101uend 777
	Water Soil Waste Oil Roste HO3 Conta	Yr Mo Dy Time	SAMPLE NOTES	EPA 502/8020 EPA 524/8240 EPA 525/8270 TLLE 22 Metals ( TPH Method- Benzene-Toluene Oil and Grease EPA 508/9080 Per
			Make Company	E23
B22 <	-1X-02201169	07001161150	Make I com posix  Analyze for  Mercony. Retain  Original sample	
B23 25		02001171235		
با			<del>}</del>	
B24 /		01:25	<u> </u>	
	Laboratory Notes :	`.	Chain of (	Custody Record
		<b>.</b>	elinquished by: (signature) Date/1	If Received by (signature )
			Inquished by: (signature ) Date/I	Ir Received by (signature )
•		₽ Pe	ilinquished by: (signature ) Date/H	Ir Received by (signature )
·		, Re	linquished by: (eignature ) Date/H	ir Received by (signature )
i		Qis	patched by: (signature ) Date/Hr	Macelved for Lab by (signature)








			Ü	han or cusive	ly rorm	, ,	ANALYSIS NECUES	IEU
	2323 Fifth S Berkeley, C	alifornia 94710	cins, Ltd		mena Jones		* ) -Xylene(s	7 )
	Matrix	Method Preserved None None None	Sample Number	Sampling Date  Yr Mo Dy Time	SAMPLE NOTES	EPA 601/8010 EPA 602/8020 EPA 624/8240	EPA 025/82/U Tirle 22 Metals EPA PP Metals ( TPH Method- Benzene-Toluene Oil and Grease	508/8080 JC 01/0
632. 35 637 44 836 44	X X X X X X X X X X X X X X X X X X X	X X X X X X X X X X X X X X X X X X X	3 1 C 1 1 9 9 0 3 2 C 1 1 9 9 0 3 3 C 1 1 9 9 0 3 1 C 1 1 9	9001190995 9001190995 900119904 9001199095 9001199095	Sance- Ho Perbov		dy Record	
				•	Relinquished by: (signature)  Relinquished by: (signature)  Relinquished by: (signature)  Relinquished by: (signature)	Date/Hr Date/Hr	Received by (s  Received by (s  Received by (s)  Received by (s)	signature )



	Curtie & Tampki		tnáin of Cu	stoc	ly Form		. ANAI	YSIS NEO	(ESIEN	(2)
	Curtis & Tompki 2323 Fifth Street Berkeley, California 94710 (415) 486-0900  Job Description W. RFS Job Number SAA-VC Client Contact	ns, Lta	Samplers	Roz	nuna Janase				\$ 523.5	
	Job Description LACRES -	- Soil begins	Phase II.					- I	`	
	Job Number 5 + A - UC.	RES-10/0			, ,			900	200	
	Client Contact Register	- Janus	Recorder	_Ra	mena Jone			(* ) 10-Xylene(s	7471	
	Matrix Method Preserved S	Sample Number	Sampling Da	θ		8010	8240 8270 Meca.	ecals od-	Grease 8080 ?	
<i>2</i> ) : 11	Warer Soll Wasce Oll Hosod Hosod None		Yr Mo Dy T	lme	SAMPLE NOTES	PA 601/	A 625/	EPA PP Mecals (* TPH Mechod-Benzene-Toluene-	A 608/	
597 1	$\times$ $\times$ $\times$ $\times$ $\times$ $\times$	1011990	9007190	115	1-3' Composite					╼╂╼╂╼
375 M		2 //	9001190	15	Jalyze for		╂	╅		
57 58	X X X	3	1001110	119						
B43 (5		2 - 2	1-1-1-2	256	11	1-1-1	- - -	- - -	- - -	
W	X	Ž /			/ /	- - -	╅	╅	-1-17	
1342 /n			3	. / 5		口				<del>        -</del>
\$ 1- \$		2 1	╂╂┼╬┼┼┼┧	<del>1</del> H	11		- - -	- - -		
				1-1-		<del> - - -</del>	┪	╂	- - -	- - -
'								世世		
	Laboratory Notes :				Chain	of Cus	lody R	ecord		
			•		elinquished by; (signature ) D	ate/l-ir		elved by	(signatur	e )
					once on mas of					
					elinquished by (signature ) D		Rec	elved by	(signatur	9 )
				Re	olinquished by: (signature ) D	ale/iler	Rec	elved by	(signatur	e )
			•	Re	elinquished by: (signature ) Di	ate/l ir	Rec	elved by	(signaturi	9)
				Ói	spatched by: (signature ) Da	11/1\el				signature )
	L						Wir	MM	<u>ulin</u>	1/19
						•		1		• '





## Curtis & Tompkins, Ltd., Analytical Laboratories, Since 1878

2323 Fifth Street, Berkeley, CA 94710, Phone (415) 486-0900

DATE RECEIVED: 12/27/89 DATE REPORTED: 02/28/90

PAGE 1 OF 4

LAB NUMBER: 19595

CLIENT: JONAS & ASSOCIATES

REPORT ON: 4 SOIL COMPOSITE SAMPLES

PROJECT #: J&A-UCRFS-101 LOCATION: UCRFS-SOIL BORING

RESULTS: SEE ATTACHED

QA/QC Officer

Laboratory Director



LABORATORY NUMBER: 19595
CLIENT: JONAS & ASSOCIATES
PROJECT #: J & A-UCRFS-101
LOCATION: UCRFS-SOIL BORING

DATE RECEIVED: 12/27/89 DATE REQUESTED: 02/15/90 DATE ANALYZED: 02/19/90 DATE REPORTED: 02/28/90

PAGE 2 OF 4

ANALYSIS: COPPER

PREPARATION METHOD: EPA 3050 METHOD REFERENCE: EPA 6010

LAB ID	SAMPLE ID	RESULT	UNITS	DETECTION LIMIT
19595-1	COMPOSITE B-18: B181122689 B182122689 B183122689	57	mg / Kg	1.0
19595-2	COMPOSITE B-12: B123122689 B122122689 B121122689	53	mg /Kg	1.0
19595-3	COMPOSITE B-38: B381011990 B382011990 B383011990	200	mg / Kg	1.0
19595-4	COMPOSITE B-39: B391011990 B392011990 B393011990	160	mg /Kg	1.0

*****************	
RPD, %	16
RECOVERY, %	77
#=====================================	



LABORATORY NUMBER: 19595
CLIENT: JONAS & ASSOCIATES
PROJECT #: J & A-UCRFS-101
LOCATION: UCRFS-SOIL BORING

DATE RECEIVED: 12/27/89
DATE REQUESTED: 02/15/90
DATE ANALYZED: 02/19/90
DATE REPORTED: 02/28/90

PAGE 3 OF 4

ANALYSIS: ZINC

PREPARATION METHOD: EPA 3050 METHOD REFERENCE: EPA 6010

LAB ID	SAMPLE ID	RESULT	UNITS	DETECTION LIMIT
	./			
19595-1	COMPOSITE B-18:	640	mg/Kg	0.50
	B181122689			
	B182122689			
	B183122689			•
19595-2	COMPOSITE B-12:	65	mg /Kg	0.50
	B123122689			
	B122122689			
	B121122689			
19595-3	COMPOSITE B-38:	170	mg/Kg	0.50
	B381011990			
	B382011990			
	B383011990			
19595-4	COMPOSITE B-39:	180	mg /Kg	0.50
	B391011990		- •	
	B392011990			
	B393011990			

RPD, %	3
RECOVERY, %	77
~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	: = = = = = = = = = = = = = = = = = = =



LABORATORY NUMBER: 19595 CLIENT: JONAS & ASSOCIATES

JOB #: J & A-UCRFS-101

LOCATION: UCRFS-SOIL BORING

DATE RECEIVED: 12/27/89
DATE REQUESTED: 02/15/90
DATE ANALYZED: 02/26/90
DATE REPORTED: 02/28/90

PAGE 4 OF 4

Analysis: Soluble Mercury

Extraction: Waste Extraction Test (WET): CCR Title 26 Section 22-66700

Analysis Method: EPA 7471

LAB ID SAMPLE ID RESULT UNITS DETECTION LIMIT

19595-1 COMPOSITE B-18: 1.3 mg/L 0.10

B181122689 B182122689 B183122689

QA/QC:

RPD, % <1
RECOVERY, % 92

Received by (signature) Received by (signature) Received by (signature) Received by (signature) **₩ ₹٢**€٥٢٧ ANALYSIS REQUESTED 908/8080 5027250 6 5C3.2 ESY and Grease Benzene-Toluene-Xylene (s) Chain of Custody Record -bodram Her PP Mecals (# E5Y Title 22 Metals 0758/8270 ESY 0128/129 Eby 0208/209 E5Y Relinquished by: (signature) Date/Hr Roman 12/27/89 Relinquished by: (signature ) Date/Hr Relinquished by: (signature ) Date/Ĥr Relinquished by: (signature ) Date/Hr Dispatched by: (signature ) Date/Hr 0108/109 ESY Make a composite to alciv. so above 30000 SAMPLE NOTES a"Fraile Fol NYYT PASE Chain of Custody Form Same Samplers Romyne Recorder Romana C 0 5 0 10 Time 0 0 Sampling Date 19 6 0 3 9 ٥ Š ; Sample Number × 20 Sor Curtis & Tompkins, Ltd Job Description UCRFS Client Contact Romana Ocher Preserved Berkeley, California 94710 None Method es: EQVH p OSS H 2323 Filth Street Laboratory Notes (415) 486-0900 \*Concainers Job Number Matrix TTO MASEE 7705 TelbW 17 <mark>ئ</mark> د Ţ

Water Client Contact Komena Job Number 34A-Job Description UCRES Laboratory Notes : Soil Matrix Waste Berkeley, California 94710 2323 Fifth Street Curtis & Tompkins, Ltd (415) 486-0900 Oil #Containers H 250 4 Preserved Method HNO<sub>3</sub> Ice None Other UCRES-Sample Number Sonar 501 Parisa d 8 Chain of Custody Form ≾ Sampling Date Z Samplers Romens Jones Recorder Roman Q Time 30 Ś Relinquished by: (signature ) Date/Hr Relinquished by: (signature ) Date/Hr Relinquished by: (signature ) Date/Hr Korneno Johan 12/27/89 Dispatched by: (signature ) Date/Hr Relinquished by: (signature ) Date/Hr ZZZIMY ZZZINI - STORY The compassion sur hun Make. I Compassite of KTEIN THE STISHED SOW FULL ANDING **SAMPLE NOTES** same da Mode above same as かいけけ かんかん Chain of Custody Record 601/8010 EPA 602/8020 EPA 624/8240 Tarenet later 625/8270 Received for Lab by ANALYSIS REQUESTED Received by (signature ) Received by (signature ) Received by (signature ) Received by (signature ) 22 Metals Metals Benzene-Toluene-Xylene(s) EPA 608/8080 Pesticides

EPA 625/8270	Dispatched by: (signature ) Date/ifr	Relinquished by: (signature ) Date/Hr	Relinquished by: (signature ) Date/Hr	Relinquished by: (signature ) Date/Hr	Romen Byar 12/27/89	Laboratory Notes : Chain of Custody			31226818912261255	X 80212268989122612:55 Some as above	841122689891122612:45	X 18 2 1 2 2 6 8 9 1 2 2 6 12:4 0 Some as above	1112268489122612:46	311226898911226112115/28250 72-9	8821226898912261215	111226848411226112115	H 2S H O I Ce Non Oth Yr Mo Dy Time	SAMPLE NOTES 601/ 602/ 624/	Method Preserved Sample Number Sampling Date	Client Contact Konsens Janes Recorder Konsens man	Job Description UCRES - Soil bolin)- Job Number S+A - UCRES - 181	Berkeley, California 94710 Samplers Nomena Jones (415) 486-0900	-
--------------	--------------------------------------	---------------------------------------	---------------------------------------	---------------------------------------	---------------------	-------------------------------------	--	--	-------------------	--------------------------------------	-----------------------	-------------------------------------------------	---------------------	----------------------------------	---------------------	-----------------------	----------------------------------------------------	-----------------------------	----------------------------------------------	---------------------------------------------------	-------------------------------------------------------------------	-----------------------------------------------------------------	---

					Laboratory Notes :		X	7 6	- 17 X 803 1 7 7 8 8 7 8 9 8 9 9 9 9 9 9 9 9 9 9 9 9		2 6 8 9 2 2 1 K 8 8 X X X X X X X X X X X X X X X X X	X	X 8121226898	8168922211118 X	H <sub>2</sub> S HNC Ice Non Oth	ntaine	Method	Romena-Jon	Job Description VCRES-Soil boning	2323 Fifth Street Berkeley, California 94710 (415) 486-0900	Tompkins, Ltd	
Dispatched by: (signature ) Date/Hr	Relinquished by: (signature ) Date/Hr	Relinquished by: (signature ) Date/Hr	Relinquished by: (signature ) Date/if	Relinquished by: (signature ) Date/Hr	Chain			Some to before	361:38	_		15	and ad	9 10 241: 115 Sames as	Yr Mo Dy Time	SAMPLE NOTES	2	Recorder Ronans Jans		Samplers Romena Jones	idii or castoug rollii	Chain of Custodu Form
Received for Lab-by (signature)	ate/Hr Received by (signature )	ate/Hr Received by (signature )	ate/itr Received by (signature )	Date/Hr Received by (signature )	Chain of Custody Record					X				E C C E	EPA	and Gr 608/80	20 40 70 [eta] als - luen ease 80 P	(# ) ne-Xyle	des 6	PCB's	ANALYSIS REQUESTED	10-7

				-	Laboratory Notes:	(61) 47) 8M 2	3	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	T	819 2	891122	 Water Soil Wast Oil #Con H2SO Ice None Other	tal	Matrix Preserved Sample Number Sampling Date	Client Contact Somma Sonab Recorder	5-101	Job Description UCKES - Soil boing	treet lifornia 900	Curtis & Tompkins, Ltd Chain of Custody F
Dispatched by: (signature ) Date/tir	Reiinquished by: (signature ) Date/Hr	Relinquished by: (signature ) Date/Hr	Relinquished by: (signature ) Date/Hr	Refinguished by: (signature ) Date/Hr Remember 12/27/89	Chain of Cu	10 ) Jume de Page		30 James Delle	_	M	5	EPA		/8010 /8020	Komena Johan	·		Comense Jonas	ody Form
(livery) deep (signature)	Received by (signature )	Chain of Custody Record			X		X	$\Box$	EPA EPA Titl EPA TPH Benz Oil	624/ 625/ e 22 PP N Meth ene- and 608/	4/8240 5/8270 22 Metals  Metals (* ) thod- e-Toluene-Xylene(s) d Grease 8/8080 Pesticides & PCB's  TUTY 7471								

	Laboratory Notes :	Curtis & Tompkins, Ltd  2323 Film Street  Berkeley, Calliornia 94710  Job Description LCRF3 - Sc., Legion  Job Number 3-1 A - UCRF5 - 1cg  Client Contact Content Sample Number Sampling Date  Preserved Sample Number Sampling Date	Thain of fustadu
Relinquished by: (signature ) Date/Hr Received by (signature )  Relinquished by: (signature ) Date/Hr Received by (signature )  Dispatched by: (signature ) Date/Hr Received for Lab by (signature )	Chain of Custody Record  Relinquished by: (signature ) Date/Hr Received by (signature )  Remained by: (signature ) Date/Hr Received by (signature )		Form

pos by rapm of Leb by (signature) Received by (signature) Received by (signature) Received by (signature) Received by (signature) MeTho Mercury ANALYSIS REQUESTED Pesticides 0808/809 ESY CLESSE pue TTO Benzene-Toluene-Xylene(s) Chain of Custody Record Method-PP Metals EBY 22 Mecals 0728/250 E5Y 078/9240 ESY Roman Jones 12/27/89 0208/209 ESY Refinquished by: (signature ) Date/Hr Refinquished by: (signature ) Date/Hr Refinquished by: (signature ) Date/Hr Relinquished by: (signature ) Date/Hr Dispatched by: (signature ) Date/Hr 0108/109 ESY Some as page ~ page 3 SAMPLE NOTES Samplers Romena Janas Chain of Custody Form Some Recorder Romens 35 Time ठ Sampling Date 3 722 72 1226 ۵ £ -68 8 9 borin) 96 Sample Number 899 Three Curtis & Tompkins, Ltd (63) Client Contact Bornelle Jouan - Soil Job Number 144 - UCRES - 10 Job Description UCRES Orper Preserved Berkeley, California 94710 (415) 486-0900 None Method Pol H 2504 Laboratory Notes 2323 Filth Street #Concainers Matrix TTO MASCE TTOS MACEE 17 12 2 ه



## Curtis & Tompkins, Ltd., Analytical Laboratories, Since 1878

2323 Fifth Street, Berkeley, CA 94710, Phone (415) 486-0900

LABORATORY NUMBER: 19172
CLIENT: JONAS AND ASSOCIATES
PROJECT #: J&A-UCRFS-101
LOCATION: UCRFS-SOIL BORING

DATE RECEIVED: 01/12/90
DATE ANALYZED: 02/01/90
DATE REPORTED: 02/02/90

5.9

======			
LAB ID	SAMPLE ID	NITRATE	SULFATE
		EPA 300.0	EPA 300.0
		mg / Kg	mg/Kg
======:			

19172-1 B211 122689/ B212 122689/ B213 122689

QA/QC:

RPD, % RECOVERY, % NITRATE 1 94 SULFATE < 1 99

QA/QC OFFICER

LABORATORY DIRECTOR

Los Angeles

Berkeley

Wilmington



## Curtis & Tompkins, Ltd., Analytical Laboratories, Since 1878

2323 Fifth Street, Berkeley, CA 94710, Phone (415) 486-0900

DATE RECEIVED: 02/21/90 DATE REPORTED: 03/05/90

PAGE 1 OF 4

LAB NUMBER: 19638

CLIENT: JONAS AND ASSOCIATES

REPORT ON: 10 SOIL SAMPLES &

4 SOIL COMPOSITE SAMPLES

PROJECT #: JA-UCRFS-101

LOCATION: UCRFS

RESULTS: SEE ATTACHED

QA/QC Officer

Laboratory Director

Los Angeles



LABORATORY NUMBER: 19638 CLIENT: JONAS & ASSOCIATES PROJECT #: JA-UCRFS-101

LOCATION: UCRFS

DATE RECEIVED: 02/21/90
DATE ANALYZED: 02/27/90
DATE REPORTED: 03/05/90

PAGE 2 OF 4

ANALYSIS: COPPER

PREPARATION METHOD: EPA 3050 METHOD REFERENCE: EPA 6010

LAB ID	SAMPLE ID	RESULT	UNITS	DETECTION LIMIT
19638-16	SS4-B18-021990	140	mg/Kg	1.0
19638-21	SS9-B39-021990	110	mg/Kg	1.0

QA/QC:

RPD, %

RECOVERY, %

108



LABORATORY NUMBER: 19638 CLIENT: JONAS & ASSOCIATES PROJECT #: JA-UCRFS-101

LOCATION: UCRFS

DATE RECEIVED: 02/21/90
DATE ANALYZED: 02/27/90
DATE REPORTED: 03/05/90

PAGE 3 OF 4

ANALYSIS: ZINC

PREPARATION METHOD: EPA 3050 METHOD REFERENCE: EPA 6010

LAB ID SAMPLE ID R	ESULT UNITS	DETECTION LIMIT
19638-16 S\$4-B18-021990	270 mg/Kg	0.50
•		
19638-21 SS9-B39-021990	150 mg/Kg	0.50

QA/QC:

RPD, % 2
RECOVERY, % 106



LABORATORY NUMBER: 19638

CLIENT: JONAS & ASSOCIATES

PROJECT #: JA-UCRFS-101

DATE RECEIVED: 02/21/90

DATE ANALYZED: 02/27/90

DATE REPORTED: 03/05/90

LOCATION: UCRFS PAGE 4 OF 4

ANALYSIS: MERCURY


PREPARATION METHOD: EPA 3020 METHOD REFERENCE: EPA 7471

LAB ID	SAMPLE ID	RESULT	UNITS	DETECTION LIMIT
19638-13	SS1-B6-021990	3.9	mg /Kg	1.0
	SS2-B3-021990	4.7	mg /Kg	1.0
19638-15	SS3-B12-021990	90	mg/Kg	10
19638-16	SS4-B18-021990	240	mg/Kg	2 0
19638-17	SS5-B16-021990	24	mg/Kg	10
19638-18	SS6-B20-021990	1.2	mg/Kg	0.10
19638-19	SS7-B10-021990	2 1	mg/Kg	10
19638-20	SS8-B37-021990	13	mg/Kg	10
19638-21	SS9-B39-021990	1.1	mg/Kg	1.0
19638-22	SS10-B43-021990	0.92	mg/Kg	0.10
19638-23	COMPOSITE B46:	14	mg/Kg	1.0
	B46-1021990			
	B46-2021990			
	B46-3021990			
19638-24	COMPOSITE B47:	4.7	mg/Kg	1.0
	B47-1021990		-	
	B47-2021990			
	B47-3021990			
19638-25	COMPOSITE B48:	0.41	mg / Kg	0.10
	B48-1021990			
	B48-2021990			
	B48-3021990			
19638-26	COMPOSITE B49:	0.32	mg/Kg	0.10
	B49-1021990			
	B49-2021990			
	B49-3021990			•

QA/QC:

RECOVERY, % 105

Laboratory Notes:	Sample Number Sampling D	
Chain of Custody Record  Relinquished by: (signature ) Date/lif  Received by (signature )  Relinquished by: (signature ) Date/lif  Relinquished by: (signature ) Date/lif  Relinquished by: (signature ) Date/lif  Received by (signature )  Dispatched by: (signature ) Date/lif  Received by (signature )  Dispatched by: (signature )	SAMPLE NOTES  SAMPLE NOTES  SAMPLE SAMPLE ABOVE  SAMPLE SAMPLE ABOVE  SAMPLE SAMPLE ABOVE  EPA 602/8020  EPA 524/9240	U J NAVITARIA SISTIFCATES SED



	JI't.	3 ().	reiller	vins,	Lite	<b>,</b>					∵.ų		7 ()			=	<u> </u>	707/	واجاد	SITEC	NES	IEU		<del></del>	
	2323 Filih Berkeley, (415) 486	Callforni	ila 94710						Sa	ımp	lers	R		Sonas 303							1.57				
	Job Descrip	otion_ <u>/</u>	ICRE:	S				_					<del></del>		11					15	1.	, I			
	Job Number				0)			•									ŀ			9					
	Client Conta					ıs		•	Re	eco	rdei	<u>_</u>	<u>.</u>	Jonas				,	2	-Xv1.	i	747/			
	· Matrix	Pi Pi	Method Preserved	Sampl	le Nu	mber		Sa	mpl	ing	Da	0		CAMPIENOTES	/8010	/ 9020	/9240	/82/0 2 Mecal	Mecais	denzane-Toluene-Xvlene(s)	and Grease	, 9080	78.		
559-B39021990	Water Soil Waste	#C30c3 #2304	HO3	_	····	<del>/</del>	Yı		للنب	Dy	1	Ime		SAMPLE NOTES	EPA 501	EPA 602	22A 524	12.8 323 11.1.0 2	22.A 29	Benzane	bus 110	M2 CC	700	2002	
	I <del>- 1 ×&gt;√</del> 1 1 1 1	-  -	<u> </u>	300	2/	790	2	<b>5</b> Ø	Z	19	0	14	4				<u>"</u>	<b>"</b>  -	╎	†"	+	TX	校	才	╁
5510.1843-021990	<del> - - - -</del>	- - -	<del>                                      </del>	3,30	4	990	1119	00	니긱	1/9	0	5	9							山		文	丁	1	
		1-1-1	╂╼╂╼╽╼╏╼╿	<u> - - -</u>	<del> - - </del>	┝┼┼	╂╌┼	+	╂	-	╂╼╂	-	H		1-1	_	4	1	_ _	11			口		
					- -		††	1	11	1	+	+-	╂┪		H	$\dashv$	+	- -	<del>                                     </del>	- -		- -	- -	- -	1
1	┟╃╂╂╂	- - -							丁	上	廿	1			1-1	+	-	-	┠╌╂╌	+	+		┟┼	╀	╀
	┟ <del>┦╌</del> ╂╌╂╌╂╌╂	-{-}-	╂═╂═┨═┦═	┠╌╁╼╂╼┦			1-1	_ _			П						1	1	<del> - -</del>	+1	-	+-1	┝╅╴	+	1-
	┟ <del>╏</del> ┼┼	╂╂╂	<del>┠╼┨╼</del> ┠╼┦	┠╾╂╼╂╼┦			╂┼	- -	4-4	- -	$\dashv$	-	$\sqcup$			$\Box$	$\perp$			口		丁		1	
		111	<del>├</del> <del>┋</del>	<u> - - -</u>	$\ \cdot\ $	┝╂┼	╂┼	+	╂┼	+	╂╾╂	-	-			4	4.	_ _	_ _	11	1				
ļ							辻	土		士						-		+	┞╌├╌	- -	- -	- -		+	
	Laboratory 1	Notes	:						,					Chain	ـــــــــــــــــــــــــــــــــــــ	~		 	 						
												1	ne	Chain elinquished by: (signature ) D			3100								
								•					2	ma mas : 2/20/90	J.Z.:	<b>45</b>		7,16	COIVE	o by	(sig	natur	• )		
			•									Re	linquished by: (signature ) Di	ate/il	r		Ne	celve	d by	(slg	natur	• )			
						Relinquished by: (signature										Date/Hr Received by (signature					• }	<del></del>			
													Re	linquished by: (signature ) D	) Date/Hr Received by (signature )										
l				*		<del></del>		<del></del>					Die	spatched by: (signature ) Da	te/l tr		V V	, n	,		/ 1	by (1		alure	· )



## Curtis & Tompkins, Ltd., Analytical Laboratories, Since 1878

2323 Fifth Street, Berkeley, CA 94710, Phone (415) 486-0900

DATE RECEIVED: 02/26/90 DATE REPORTED: 03/08/90

PAGE 1 OF 4

LAB NUMBER: 19697

CLIENT: JONAS & ASSOCIATES

REPORT ON: 4 SOIL COMPOSITES &

1 WATER SAMPLE

PROJECT #: JA-UCRFS-101

LOCATION: UCRFS

RESULTS: SEE ATTACHED

QA/QC Officer

Laboratory Director

Berkeley Wilmington Los Angeles



LABORATORY NUMBER: 19697 CLIENT: JONAS & ASSOCIATES PROJECT #: JA-UCRFS-101

LOCATION: UCRFS

DATE RECEIVED: 02/26/90 DATE ANALYZED: 03/06/90 DATE REPORTED: 03/08/90

PAGE 3 OF 4

ANALYSIS: MERCURY

METHOD REFERENCE: EPA 7470

LAB ID SAMPLE ID RESULT UNITS DETECTION LIMIT

19697-13 MW122590 ND mg/L 0.001

ND = NONE DETECTED

QA/QC Summary:

RPD, % 2
RECOVERY, % 113



LABORATORY NUMBER: 19697 CLIENT: JONAS & ASSOCIATES PROJECT #: JA-UCRFS-101

LOCATION: UCRFS

DATE RECEIVED: 02/26/90
DATE ANALYZED: 03/02/90
DATE REPORTED: 03/08/90

PAGE 2 OF 4

ANALYSIS: MERCURY

METHOD REFERENCE: EPA 7471

LAB ID	SAMPLE ID	RESULT	UNITS	DETECTION LIMIT
19697- 1,2,3	MWB1322490/ MWB12.522490/ MWB24.522490	36	mg /Kg	10
19697 - 4,5,6	MWB3522490/ MWB35.522490/ MWB3622490	27	mg / K g	10 .
19697- 7,8,9	MWB 4 6 . 5 2 2 4 9 0 / MWB 4 7 2 2 4 9 0 / MWB 4 7 . 5 2 2 4 9 0	2 4	mg / Kg	10
19697 - 10,11,12	MWB5822490/ 2 MWB58.522490/ MWB5922490	2.7	mg/Kg	1.0

O۷	4/	$\mathbf{OC}$	S	u mm a	r	ν	:
----	----	---------------	---	--------	---	---	---

RPD, %	8
RECOVERY, %	106



LABORATORY NUMBER: 19697 CLIENT: JONAS & ASSOCIATES PROJECT #: JA-UCRFS-101

LOCATION: UCRFS

19697-13 MW122590

DATE RECEIVED: 02/26/90 DATE ANALYZED: 03/02/90 DATE REPORTED: 03/08/90

25

PAGE 4 OF 4

ANALYSIS: TOTAL DISSOLVED SOLIDS

METHOD REFERENCE: EPA 160.1

mg/L

LAB ID SAMPLE ID RESULT UNITS DETECTION LIMIT

1,300

		•	'	】 しし 232	Jr 3 F	LTS Tith	SI	991	1 (	n C	ip	K	īn	\$	,	L t	d		(			انبا	<b>)</b>		•			F-1	) (	Jones 161/	F		_   	<del>,,</del>	M	.YSK	SNE	П	П	U					
			L	(41	5)	486	-09	00		947								_1					S	am	pl	ers	3_	K	. (	Jones									s, EDé				,		
(	93)	J	ob	D	esc	crip	lio	n_	_6	10	_/2	2 F	:3														***								١		(5)		3 5	1					
( )	< 5 <sub>/</sub>	) J	ob	Nu	mb	eĽ	3	EZ	<del>}_</del> -	- L	<u>C</u> ,	12	E	5	_	10	رن												1	^	1				-	_	-Xylene(s)		0.00	7					
		C	lier	nt	Co	nta	act	<u> </u>	K	M	ىقا	N	<u>ب</u>	2		no	ine.	2					F	lec	oro	de 	r_	ŀ	7	Jonas					51	3	X-90			飞					
		Ŀ	· N	/lai	rix	<del></del>	2023	Ŀ		thod 9 0 1 0 5			Sa	ım	ple	N	un	ıbe	er			Sa	mp	oling	g (	Da	te				8010	8020	9240	8270	Meta	ecals	To Lue	Greas	9080	2702					
		1097	Soil	1358	!		#Concai	304	HQ3	10.0	ge .	ner								L		-		<del></del>	-1				_	SAMPLE NOTES	601/	602/	524/	625/	.e 22	D: 0:	Respondent de	puz	908	2					
	,	× ×	Š	3	ö	+	15	3:	f	H		<u> </u>	14			_	1.	T <sub>G</sub>		٠.	Yr		7	D	y		in	18			EPA	Si	EPA	EZA	;;	EP.A	Ben H	017	E23	7					
	_	-	X	$\dashv$	╅	╁	╢	<del> </del>	-	$\bowtie$	- -	- -	3 6		3 2	- 12		45	10		10	10	2	2	4	a	9	4	9	For MITCUTY.	_	L				$\Box$	I			$\nabla$	$oldsymbol{\mathbb{T}}$	П	_		
c I	7	-	H	1		╁	╁	-		r		+	14	<u>'</u> h	4	╁	╂	4-	1-	-{ -	╁	╢	╂┈	14	4	-	-	$\dashv$		Retain original Sample	-	<b> </b> _	_	-		$\dashv$	- -	╄	11	4	_ _	$\sqcup$			
	3		X	176745													╁	칻	扌	†	$\dagger$	十	1	H	1	ō	9	3	0		-	-	-	H	$\dashv$	+	- -	╀	H	4	- -	╀	_		
	.,	. _																									$\Box$					1-	-			+		十	╁╅	4	+-	╁┼	-		
01	, T	-	$\stackrel{\bigcirc}{\times}$	-	4	- -	- -	<u> </u>	_	X	_ -				<u> </u>	2 -	45	42		9	10		7	2	4	0	2	3	3	LANGE FUL						十	1	<u> </u>	$\Box$	J	1	H	_		
٠. ر	6	$\vdash$	$\Theta$	$\dashv$	+	╬	┨–	╁	-	X			1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1	_		42	1	1/2	4-	-1-4			2		4				2	Mercury. Retain									口	X			_		
	•	$\vdash$	H		+	╁	╁	╂─	-	$^{H}$	+	- [	, T	4	2	-   -	17	45			10	10	Z	. 2	4	Q	4	3	<u>5</u> ]	About Sofuture	_	_	_			4	_ _	L		$\Box$	$oldsymbol{\square}$	$\prod$	_		
					+	+	╁	╁			- -	- -	-{-	+	╁	╁	╁	十	╁╴	╁	╁	╁	╂╌	$\vdash$	$\dashv$	$\dashv$	-	-			-	┡	_	-	4	4	4	<del> </del>	_	4	_ _	14	_		
								匚			1	土	土	1	上	1	土	İ	土	1	土	上						_	-		-	-	-	╂┤	$\dashv$	-	+	╁	╂╌╂	+		╟	_		
		Tu	.ab	ora	tor	y 1	Vol	es	:			<del></del>				•							`					1		Chain	ol	Cu	sto	odv	R		ord						_		
																													R	elinquished by: (signature ) D			T				ed by	y (!	slgna	alure	<u> </u>		_		
																						•							R	man 10000 2/26/90	7 3	: Z	۵				•	•	•		•				
																														elinquished by: (signature ) D					Rec	:elve	ed by	/ (8	ilgna	iture	)				
																													Re	elinquished by: (signature ) D	ate/l	Ir		1	Rec	:elve	rd by	r (9	ilgnature )						
																													Re	elinquished by: (eignature ) D	ate/l	Ir	1		Rec	:elve	ed by	/ (:	ngna	ilur e	)		-		
																													Dİ	spatched by: (signature ) Da		ir	-	4	Nec	elve	To the	<u></u>			Ignati	lure)			
															-							<del></del>						I					7	<	1	t a	11	X		<u>ک د</u> ۱	<u>~</u>	<del>/</del> 20			
																																			1		V			1	٠, س				

•		$\overline{}$	ur	Ti.	\ {	7		m	ħ	Z i	n		ı	7	_	ì	-	_	•			•			L	, =			_	·	170	W/L	า SIS	HEC	<b>WES</b>	STEL			-	
		1 2	2323 Berkel 415)	Filih øy, (	Str Cali	eet forni			_			',	L. \							S	am	ıpl	ler	s_	R		Jonas									2C3.				
		Job	Des	crip	lio	n	U	(	RI	= 3	>													-										Î	1.	J.				
		Job 1	lmuV	ber		<u>S</u> /	<del>1</del> -	- ر	) C	R	E:	5	<u> </u>	10	ٔ د									-										2						
		Clien	t C	onla	act	<u> </u>	<u> </u>	571	ce,	nu	<u>_</u>	Ć	/	377	a	<del>6</del>				F	lec	or	de	r	R	•	Jonas					2	١	ne-Xvlene(s)		7950101des	7			
		. · M	latrix	: 	ners	l	Meth	bor 1	<u>d</u>	s	an	ple	1 e	lun	nb	er		;	Sa	mp	lin	g	Da	ıle				601/8010	8020	8240	8270	Meta	ecals	Tolue	Greas	9080				
		Water	11		Concai	2304	ဌ	o Ce	Cher								-		1.		Τ.,		Γ.				SAMPLE NOTES	•	4 602/	1 524/	625/	e 22	EPA 39 Mecals	-aua21	Oil and Grease	/809	30			
	7	3 0	* 0	X x x x 2 2 4 9 0 90 0 2 2 4 0 9 4 C														_	7	EPA	品	E2?	E23		il E	36	9	52A												
C3	8	TXT	11	Yr Mo Dy Time  Yr Mo Dy Time  Yr Mo Dy Time  X														Mercury Patrice	-	_	-		4-	- -	4-1	_	4_	1	Ц	$\perp$										
	Ģ	X																Sample for FUTURE	-	╂╌	-	$\vdash$	+	十	H	┟╌┨	$-\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!$	<b>\</b>  _	╂╌╂╌	╀										
		-	44	_ _	lacksquare		_			. _		. 1		ı		1	ı	1	ı			i					7 : 10/4/73/73		-	-	$\sqcap$	十	- -	$\dagger \dashv$	┟─╂	-	╢	╁┼	╂	
64	/•	+	X 3 8 2 2 4 9 0 90 0 2 2 4 0 9 5 0																			1				十	1	<del>                                      </del>	十											
CT	12		+		╁	H	-k	<del>X</del>  -	╁	1,3	87 87	3	-	+	4	4-	-	-	+	栏	1-1	_	Н	4	$\dashv$	_	) Same as above			Ц		$\perp$		$\square$	$\Box$	$\Box$		П	I	
	1 6.		11	$\top$	┪	$\Box$	ť	4-	╁	H	<del>2</del> 4	4	+	╁	f	7	- -	+-	╁	F	H	Н	$\vdash$	4	-	-	/	-	_			4	4	$\sqcup$	$\sqcup$		1		$\perp$	
									1	17	7	+	+	†	$\dagger$	╁	╁	╁╴	╁	十	H	$\vdash$	$\vdash$	+	-	-		-		Н	$\vdash$	+	╀	╂╂	┟╼╂	- -	╬	╀	╀	
		<b> - - </b>	- -	_	1_	П	$\perp$																					-	1-		┟┼	+	十	H		┰	╁	╂╌╂╌	╁	
			<u> </u>		<u></u>	Ш							_L		_L					L									上		口	土	土	廿		土	$\pm$		+	
		Labo	rato	ry N	Vot	es	:													`							Chain	of	Cu	slo	-dy	Re	 BCO	rd						
																			•								elinquished by: (elgnature ) D			T	ī	Rec	elve	d by	(sk	gnat	UTB	)		
		İ																								IV.	ome 2/26/9 elinquished by: (signature ) D	60	<u>۲: ی</u>	4-	<del>_</del>					<del></del>				
																																190	91V 8(	d by	(3)	gnati	XO )	)		
																										Re	elinquished by: (signature ) D	ate/l	lr.		F	lec	əlvər	d by	(sle	gnati	NO 1	)		
																										Re	elinquished by: (signature ) D	ale/l	Hr	1	-	lec	elve:	d by	(a)	gnati	NO	)		
																										DI	spatched by: (signature ) Da	te/1 i	r	T	1	Deceived to Lab by (signature 2-26-90)								
																				<del></del>					J L	L					1	<u> </u>	1	1/1	7	<u> </u>	<u>ئے۔</u> ا	<u>; 2</u>	. 0	

Charles 100

**-**

A 25.

Merculy Received by (signature) Received by (alguature) Received by (signature) Received by (signature) <u>50</u> . MYNLTSIS RECUESTED \$,ED6 9 \$80;0;3886 0808/809 453 and Grease Senzene-Toluene-Xylene(s) Chain of Custody Record -bodrank Rer 4) STEDEN GC ¥23 TIETA ZZ Mecals 075/8270 078/729 ESY 1700 2/2/Ph 3:2 0208/209 EBY Refinquished by: (signature ) Date/14r Relinquished by: (signature ) Date/it Relinquished by: (signature ) Date/1 in Relinquished by: (signature ) Date/Hit Dispatched by: (signature ) Date/Hr 0108/109 ESY SAMPLE NOTES Committee Committee Committee Rome 0205 Samplers K. Recorder R. Time Sampling Date 25 ò 20 £ 90 ۲ 22 590 Sample Number curtis & iompkins, Ltd 000 <u>ड</u> ह Client Contact Ronwa Job Number JA - UCR FS Job Description <u>UCRES</u> 29230 2323 Flith Street Berkeley, California 94710 (415) 486-0900 anov. Method eo] HO3 Laboratory Notes: \$ 05g H \*Concainers Matrix TTO Masce TTOS Waces